


MERCK

REFINE PROTEIN PREPARATION.

Tools for better protein analysis.

The life science business of Merck operates as MilliporeSigma in the U.S. and Canada.



Sigma-Aldrich[®]
Lab & Production Materials

Millipore[®]
Preparation, Separation,
Filtration & Monitoring Products

INTRODUCTION

Today, researchers are challenged to create high quality samples for meaningful protein analysis, often using cumbersome traditional sample preparation methods. With over 50 years of experience in developing protein sample preparation technologies, Merck is constantly innovating new tools to offer you rapid and efficient solutions that can be smoothly integrated into your workflow.

Why spend your time on arduous sample preparation protocols when you can focus your efforts on exciting experiments? With the right pure protein, in the buffer you need, at the concentration you want, your next discovery is only a step away. From protein isolation to purification, you can count on us to support your research with maximum yields of intact, functional proteins.

To learn more, please visit: SigmaAldrich.com

KEY FEATURES

Unmatched Flexibility

Isolate proteins from a diverse range of sample types with our flexible, broad range of kits.

Diverse downstream applications

Our reagents enable you to produce samples that can be used directly in applications such as activity assays, protein microarrays, SDS-PAGE, immunoblotting, ELISA, two-dimensional gel electrophoresis (2DGE), mass spectrometry (MS; including MS/MS, LC-MS, MALDI-MS, SELDI-MS, and ESI-MS), and others.

Scale-up compatibility

It's easy to scale up to high-throughput recombinant protein purification and solubility screening using our sample preparation reagents.

Sigma-Aldrich®

Merck offers, under the Sigma-Aldrich® brand, a strong and always-expanding portfolio of lab and production materials that keep our customers' important work moving forward. And through our technical support and our scientific partnerships, we help connect our customers to a whole world of progress.

 Denotes Sigma-Aldrich® Products

Millipore®

Merck offers, under the Millipore® brand, an ecosystem of industry-leading products and services, spanning preparation, separation, filtration and monitoring — all of which are deeply rooted in quality, reliability and time-tested processes.

 Denotes Millipore® Products

TABLE OF CONTENTS

| | |
|---|-----------|
| Protein Extraction | 4 |
| Protein Extraction with Cell Lysis Reagents (“Busters”) | 5 |
| Fast, simple, gentle protein extraction from <i>E. coli</i> , yeast, insect and mammalian cells | |
| Cell Lysis and Nucleic Acid Removal Enhancers | 7 |
| Increase protein extraction yield and purity with Benzonase® and rLysozyme™ reagents | |
| Protein Extraction with ProteoExtract® Kits | 9 |
| Extract proteins from different fractions of the mammalian cell, including membrane, nucleus, cytosol and cytoskeleton | |
| Protein Extraction with Inhibitors | 11 |
| Protease and phosphatase inhibitor cocktails to prevent proteolysis and dephosphorylation of proteins | |
| Protein Purification and Depletion | 13 |
| Affinity Purification with PureProteome™ Magnetic Beads | 14 |
| Ideal for small volume applications such as immunoprecipitation and recombinant protein screening | |
| Affinity Purification with Recombinant Fusion Tags | 17 |
| Protein Depletion | 22 |
| Protein depletion reagents: Seppro® Protein Depletion, PureProteome™ reagents, ProteoExtract® depletion, ProteoPrep® products | |
| Amicon® Pro Purification System | 24 |
| Purify, exchange buffer and/or concentrate in a single device | |
| Protease Cleavage Enzymes. | 26 |
| Recombinant enterokinase, factor Xa, HRV 3C protease, thrombin and other enzymes for cleaving fusion proteins | |
| Protein Buffer Optimization and Sample Concentration | 28 |
| Dialysis and Buffer Exchange Devices | 29 |
| Amicon® Pro Purification System, D-Tube™ Dialyzers and Amicon® Ultra centrifugal filters for protein sample diafiltration and buffer exchange | |
| Centrifugal Concentration Devices | 32 |
| Amicon® Ultra filters for fast and effective protein concentration | |
| Specialized Concentration Devices | 37 |
| Microcon®, Ultrafree® and Centriprep® filters for efficient purification, concentration, and desalting of biological samples | |
| Clinical Filtration Devices. | 40 |
| Centrifree® and Minicon® concentrators for concentration or partition of body fluids or other biological specimens | |
| Large Volume Concentration Devices | 41 |
| Centricon® Plus-70 centrifugal filters for rapid processing of aqueous biological solutions in larger volumes; stirred cells and cut discs for concentrating volumes up to 400 mL | |

PROTEIN EXTRACTION

When purifying proteins for functional or structural studies, the first step is to disrupt the cells or tissue sample and extract the relevant protein fraction. This step is critical because processing methods that require harsh mechanical, chemical, or enzymatic treatments can affect the target protein's integrity and activity, or otherwise expose it to degradative conditions.

Our complete range of reagents and enzymes for cell lysis and protein extraction provide you with an array of options so that you can put together the perfect extraction protocol for your particular cells and protein.

Protein Extraction Reagents Application Guide

| Products by Cell Type | Starting Material | | | Applications | | | Comments |
|--|-------------------|-------------|--------|--------------|---------------|----------------|--|
| | Total Culture | Cell Pellet | Tissue | 1D PAGE | 2D PAGE / IEF | Activity Assay | |
| <i>E. coli</i> | | | | | | | |
| BugBuster® Master Mix | | ■ | | ■ | ■ | ■ | Combines BugBuster® Protein Extraction Reagent with Benzonase® Nuclease and rLysozyme™ Solution. Convenient, all-in-one protein extraction reagent efficiently lyses bacteria and digests nucleic acids. |
| BugBuster® Protein Extraction Reagent | | ■ | | ■ | ■ | ■ | Efficient protein extraction from <i>E. coli</i> under non-denaturing conditions. |
| BugBuster® 10X Protein Extraction Reagent | | ■ | | ■ | ■ | ■ | A concentrated form of BugBuster® Protein Extraction Reagent. Ideal for extraction when a specific buffer is required for protein stability. |
| PopCulture® Reagent | ■ | | | ■ | | ■ | Protein extraction from cells directly in the culture medium; no centrifugation required. |
| Yeast | | | | | | | |
| YeastBuster™ Protein Extraction Reagent | | ■ | | ■ | | ■ | Efficient protein extraction from yeast under non-denaturing conditions from any volume of culture. Add 0.5 M THP Solution (included) and Benzonase® Nuclease for enhanced efficiency. |
| Insect | | | | | | | |
| CytoBuster™ Protein Extraction Reagent | | ■ | | ■ | ■* | ■ | Gentle lysis of insect cells with retention of protein activity for assays and purification. Can use with monolayers or pellets derived from suspension cultures. |
| Insect PopCulture® Reagent | ■ | | | ■ | | ■ | Lysis of insect cells directly in serum-free medium. Ideal for expression screening of many small samples. |
| Mammalian | | | | | | | |
| CytoBuster™ Protein Extraction Reagent | | ■ | | ■ | ■* | ■ | Gentle lysis of mammalian cells with retention of protein activity for assays and purification. Can use with monolayers or pellets derived from suspension cultures. |
| ProteoExtract® Kits | | ■ | | ■ | ■* | ■ | Extract protein fractions from different parts of the cell. A range of kits offering maximum flexibility. |
| Stabiliser™ Reagent | | | ■ | ■ | ■* | ■ | Stabilizes functional protein and maintains nucleic acid integrity. Prevents degradation during tissue lysis and storage. |
| Lysis and Extraction Enhancement | | | | | | | |
| Gram-negative bacteria (<i>E. coli</i>) | | | | | | | |
| rLysozyme™ Solution | ■ | ■ | | ■ | | ■ | Cleaves bond in peptidoglycan layer of <i>E. coli</i> cell wall. |
| Lysonase™ Bio-processing Reagent | ■ | ■ | | ■ | | ■ | Convenient mixture of rLysozyme™ solution and Benzonase® Nuclease minimizes pipetting steps. |
| Gram-positive bacteria | | | | | | | |
| Chicken Egg White Lysozyme Solution | ■ | ■ | | ■ | | ■ | Cleaves bond in peptidoglycan layer of bacterial cell wall. |
| All cells | | | | | | | |
| Benzonase® Nuclease | ■ | ■ | | ■ | | ■ | Degrades all types of nucleic acids for more efficient protein extraction, faster chromatography, and reduced interference in assays. |

1D PAGE — One-dimensional Polyacrylamide Gel Electrophoresis

2D PAGE — Two-dimensional Polyacrylamide Gel Electrophoresis

IEF — Isoelectric Focusing

* — Salt must be removed before IEF

Protein Extraction with Cell Lysis Reagents (“Busters”)

Featured Products


Stabiliser™ Reagent

DNA, RNA and protein stabilization in tissue extracts


Maintain nucleic acids and functional proteins with Stabiliser™ reagent, the ONLY reagent that protects the integrity of nucleic acids AND active protein in one uniform lysate mixture. Now, you can stabilize complete, uniform tissue homogenates providing sample-to-sample comparability. The convenient all-in-one Stabiliser™ formulation protects analytes at the time of lysis and provides protection from freeze/thaw cycles during long-term storage for future detection needs:

- Maintain functionally active protein and nucleic acids from the same tissue samples
- Long-term storage and protection from freeze/thaw cycles for future detection needs
- Archive tissue samples if future analyte needs change

A.



B.




Stabiliser™ reagent extracts higher yields of protein, ~5X more than PBS buffer and 2X more than CytoBuster™ reagent. Additionally, Stabiliser™ reagent protects enzymatic activity during multiple freeze/thaw cycles. (A) 250 mg sections of chicken hearts were lysed in 5 mL CytoBuster™, Stabiliser™ reagents or PBS buffer. Samples were normalized and loaded at 10 µg protein per lane. (B) Enzymatic activity was measured every two days after a freeze/thaw cycle using our IDH activity assay kit on samples stored at -20 °C.

BugBuster® Protein Extraction Kits and Reagents


Simple extraction of soluble protein from *E. coli*, without sonication

Gently disrupt the cell wall of *E. coli* and liberate soluble proteins with BugBuster® Kits and Reagents. BugBuster® reagent provides a simple, rapid, low-cost alternative to mechanical methods such as French press or sonication for releasing expressed target proteins in preparation for purification or other applications. The proprietary formulation uses a detergent mix to perforate cell walls without denaturing soluble protein. Simply harvest cells by centrifugation and suspend in BugBuster® reagent. Following a brief incubation, remove insoluble cell debris by centrifugation. The clarified extract is ready to be purified.

A.



B.



BugBuster® reagent is superior to “homebrew” lysis buffer and BugBuster® reagent with both Benzonase® nuclease and rLysozyme™ solution produced lysates with the highest 6XHIS-CRP yields. (A) *E. coli* lysates (5 µL of 1 mL total lysate) from various lysis protocols were fractionated and analyzed by SDS-PAGE. A band corresponding to 6XHIS-CRP is prominently visualized in the BB +/+ lane. (B) Cleared cell lysates (2 µL of 1 mL total) were spotted on assay cards and quantified using the Direct Detect® spectrometer. In each case, bars represent the average of 3 independent samples.

How do I choose between BugBuster® Products?

Components of Bacterial Lysis Reagents

| | BugBuster® Reagent | Buffer | Benzonase® Nuclease | rLysozyme™ Solution | Notes |
|--|--------------------|--------|---------------------|---------------------|--|
| BugBuster® Reagent | ■ | ■ | | | |
| BugBuster® 10X reagent | ■ | | | | Flexibility to customize dilution and buffer composition |
| BugBuster® Plus Benzonase® Nuclease | ■ | ■ | ■ | | 2 separate vials for greater flexibility |
| BugBuster® Plus Lysonase™ Kit | ■ | ■ | ■ | ■ | 2 separate vials for greater flexibility |
| BugBuster® Master Mix | ■ | ■ | ■ | ■ | 1 convenient reagent |
| PopCulture® Reagent | ■ | ■ | | | Buffer protects protein from the pH extremes produced in high-density culture media, enabling extraction directly in medium. |

We offer a family of protein extraction reagents for gentle, efficient, non-mechanical extraction of soluble proteins from bacteria, yeast, plant, mammalian, and insect cells. ●

CytoBuster™ reagent — Obtain protein extracts from mammalian and insect cells in their native state, in 5 minutes.

NucBuster™ reagent — Extract nuclear proteins in less than 30 minutes with a simple 2-step protocol.

PhosphoSafe™ Extraction reagent — The PhosphoSafe™ Extraction Buffer is a detergent and phosphatase inhibitor mixture optimized for fast, efficient extraction of soluble proteins from mammalian and insect cells that preserves the phosphorylation state of sample proteins.

YeastBuster™ reagent — Extract proteins from yeast and plants without mechanical disruption or enzymatic lysis. The reagent has been tested with

Saccharomyces cerevisiae, *Pichia pastoris*, *P. stipidis*, and *Schizosaccharomyces pombe* strains, and with plant cells.

Insect PopCulture® reagent — Insect PopCulture® Reagent is a detergent-based lysis reagent specifically formulated for extraction from total insect cell culture (in suspension or adherent) without the need for centrifugation.

Stabilyser™ reagent — Use Stabilyser™ reagent and protect both nucleic acids and functional proteins in one uniform lysate mixture. Stabilyser™ reagent provides long-term storage and protection from freeze/thaw cycles.

Ordering Information

| Application | Description | Catalog No. |
|-------------|---|-------------|
| Bacteria | BugBuster® Protein Extraction Reagent | 70584 |
| | BugBuster® Master Mix | 71456 |
| | BugBuster® Plus Benzonase® Nuclease | 70750 |
| | BugBuster® Plus Lysonase™ Kit | 71370 |
| | BugBuster® 10X Protein Extraction Reagent | 70921 |
| | PopCulture® Reagent | 71092 |
| Mammalian | CytoBuster™ Protein Extraction Reagent | 71009 |
| | NucBuster™ Protein Extraction Reagent | 71183 |
| | PhosphoSafe™ Extraction Reagent | 71296 |
| | Stabilyser™ Reagent | PNS1010 |
| Yeast | YeastBuster™ Protein Extraction Reagent | 71186 |
| Insect | Insect PopCulture® Reagent | 71187 |

CellLytic™ Lysis Reagents ►

Features and Benefits

- **Efficient:** Higher protein extraction efficiency than traditional methods such as sonication and lysozyme
- **Non-denaturing:** Does not interfere in downstream applications including immunoprecipitation, kinase and phosphatase assays, reporter gene assays and gel shift assays
- **Convenient, ready-to-use reagent**

CellLytic™ reagents are specifically formulated to lyse and extract cellular proteins based on the type of expression system. All are designed to rapidly lyse the

cells with an easy-to-follow protocol. The CellLytic™ family is compatible with a wide variety of protease inhibitors, chelating agents, and chaotropes. Because the proteins are in a non-denaturing environment, these reagents do not interfere with standard affinity chromatography. Downstream applications, such as Western blots, gel-shift assays, affinity purification, and reporter detection can be performed without removing the CellLytic™ reagent. Overall extraction efficiency is consistently higher than with other common protocols, such as freeze-thaw or sonications. Although each CellLytic™ reagent is uniquely formulated, all are amenable to scale-up to meet any laboratory's needs.

Ordering Information

| Application | Description | Catalog No. |
|-------------|---|----------------|
| Bacteria | CellLytic™ B Cell Lysis Reagent, 10X Concentrate | C8740 |
| | CellLytic™ B Cell Lysis Reagent, 2X Concentrate | B7310 |
| | CellLytic™ B Cell Lysis Reagent, Standard Strength | B7435 |
| | CellLytic™ B Plus Kit | CB0050, CB0500 |
| | CellLytic™ Express reagent, for in-culture bacterial cell lysis | C1990 |
| | CellLytic™ Express reagent, 1 mL tablets, for direct lysis of bacterial cultures and for use in the His-Select® iLAP column | C5236 |
| Mammalian | CellLytic™ M, Cell Lysis Reagent | C2978 |
| | CellLytic™ Mem Protein Extraction Kit, for membrane proteins | CE0050 |
| | CellLytic™ MT Cell Lysis Reagent, for mammalian tissues | C3228 |
| | CellLytic™ Nuclear™ Extraction Kit, for mammalian tissue or cultured cells | NXTRACT |
| Plant | CellLytic™ P Cell Lysis Reagent | C2360 |
| | CellLytic™ PN Isolation/Extraction Kit, for plant leaves | CELLYTPN1 |
| Yeast | CellLytic™ Y Cell Lysis Reagent | C4482 |
| | CellLytic™ Y Plus Kit for enzymatic yeast cell lysis | CYP1 |

Cell Lysis and Nucleic Acid Removal Enhancers

Featured Products

Benzonase® Nuclease ●

Effectively reduce viscosity and remove nucleic acids from protein solutions

Benzonase® Nuclease is a genetically engineered endonuclease from *Serratia marcescens*. It degrades all forms of DNA and RNA (single stranded, double stranded, linear and circular) while having no proteolytic activity. It is effective over a wide range of conditions and has an exceptionally high specific activity. Benzonase® nuclease is an excellent choice for viscosity reduction to shorten processing time and increase protein yields.

Benzonase® Advantages

- Compliant with FDA guidelines for nucleic acid contamination
- Functional between pH 6 and 10, from 0 °C to 42 °C, for maximum versatility
- Active in the presence of ionic and non-ionic detergents, reducing agents, PMSF (1 mM), EDTA (1 mM) and urea.
- Available in ultrapure (> 99% by SDS-PAGE) and pure (> 90%) grades
- Available in standard concentration (25 U/μL) and high concentration (HC, 250 U/μL).



Nucleic acid digestion by Benzonase® Nuclease.
E. coli BL21(DE3) cells containing a pET construct were suspended in BugBuster® Reagent (5 mL/g wet weight). Identical volumes of the suspension were treated with the indicated amounts of Benzonase® Nuclease for 30 min at room temperature. Samples were clarified by centrifugation and analyzed by agarose gel electrophoresis and ethidium bromide staining.

***E. coli* lysate without Benzonase® Nuclease.**
 Gooney, viscous, difficult to handle.

rLysozyme™ Solution

Degrade bacterial cell walls with stabilized recombinant lysozyme

rLysozyme™ Solution contains a highly purified and stabilized recombinant lysozyme that can be used for lysis of *E. coli*. The enzyme catalyzes the hydrolysis of N-acetylmuramide linkages in bacterial cell walls. The specific activity of rLysozyme™ solution (1700 KU/mg) for *E. coli* lysis is 250 times greater than that of traditional chicken egg white lysozyme. rLysozyme™ solution is optimally active at physiological pH. Very small amounts of rLysozyme™ solution enhance the efficiency of protein extraction with BugBuster® and PopCulture® Reagents. The product is supplied as a ready-to-use solution and is stable at -20 °C.



rLysozyme™ solution exhibits 250 times higher specific activity than chicken egg white activity when measured using a standard activity assay.

Ordering Information

| Description | Catalog No. |
|--------------------------------------|-------------|
| Benzonase® Nuclease, Purity > 90% | 70746 |
| Benzonase® Nuclease HC, Purity > 90% | 71205 |
| Benzonase® Nuclease, Purity > 99% | 70664 |
| Benzonase® Nuclease HC, Purity > 99% | 71206 |
| rLysozyme™ Solution | 71110 |
| Chicken Egg White Lysozyme Solution | 71412 |
| Lysonase™ Bioprocessing Reagent | 71230 |

Protein Extraction with ProteoExtract® Kits

Featured Products


ProteoExtract® Subcellular Proteome Extraction Kit (S-PEK) ●

Reproducible extraction of subcellular proteomes from mammalian cells.

Based on different solubilities of certain subcellular compartments, the S-PEK uses proprietary chemistries to yield four subproteome fractions which are enriched in cytosolic, membrane/organelle, nuclear, and cytoskeletal proteins. In the case of adherent cells, the procedure is performed directly in the tissue culture dish without the need for cell removal. For suspension-grown cells, extraction starts with gentle sedimentation and washing of cells. Extraction from tissues requires isolation of viable cells before proceeding with the extraction protocol.

Applications of S-PEK:

- Subcellular redistribution assays to monitor protein translocation
- Enzyme activity assays including reporter gene assays and kinase assays
- SELDI (surface-enhanced laser desorption/ionization) profiling
- Non-denaturing gel electrophoresis
- Assaying protein expression levels using fluorescently-labeled subcellular extracts in microarrays




ProteoExtract® Native Membrane Protein Extraction Kit (M-PEK)

Isolation of native membrane proteins from mammalian cells and tissue.

Extract proteins associated with cellular membranes with M-PEK. Extremely mild extraction conditions yield a 3–5 fold enrichment of integral membrane and membrane-associated proteins. The simple, two-step procedure enables processing of multiple samples in parallel. Extraction from tissues requires isolation of viable cells before proceeding with the extraction protocol.

Applications for Extracted Membrane Proteins:

- Enzyme activity assays, including reporter gene assays and kinase assays
- Non-denaturing and denaturing gel electrophoresis, immunoblots and immunoassays
- Assaying post-translational modifications, such as phosphorylation
- SELDI-profiling of integral and membrane-associated proteins
- NHS ester labeling of membrane proteins



Notably increased enrichment of EGF receptor using M-PEK compared to total cell lysate. HEK293 cells were extracted with buffered 1% Triton® X-100 surfactant to generate a total lysate or extracted with M-PEK to yield a membrane fraction. Equal volumes of these fractions were utilized to quantitate the concentration of EGF receptor in the samples using an EGF-R ELISA Kit. Protein concentrations were used to calculate the amount of EGF-R per mg protein in the total lysate and the membrane fraction. The measurements demonstrate a 4.5 fold enrichment of the EGF receptor in the M-PEK-extracted membrane fraction.

Ordering Information

| Application | Description | Catalog No. |
|------------------------------|---|-------------|
| Organelle Fractionation | ProteoExtract® Subcellular Protein Extraction Kit | 539790 |
| | ProteoExtract® Complete Mammalian Protein Extraction Kit | 539779 |
| | ProteoExtract® Cytosol/Mitochondria Fractionation Kit | QIA88 |
| | ProteoExtract® Native Cytoskeleton Enrichment Kit | 17-10210 |
| | ProteoExtract® Cytoskeleton Enrichment and Isolation Kit | 17-10195 |
| Membrane Proteins | ProteoExtract® Native Membrane Protein Extraction Kit | 444810 |
| | ProteoExtract® Transmembrane Protein Extraction Kit | 71772 |
| Mass Spec Peptide Enrichment | ProteoExtract® All-in-One Trypsin Digestion Kit | 650212 |
| | ProteoExtract® Glycopeptide Enrichment Kit | 72103 |
| | ProteoExtract® Phosphopeptide Enrichment TiO ₂ Kit | 539722 |

ProteoPrep® Lysis Kits

ProteoPrep® kits and individual extraction reagents allow for selective or total protein extracts from cellular samples. The protein extractions obtained with each component can be optimized to meet your individual needs. The reducing and alkylating reagents produce protein samples that exhibit improved focusing and decreased streaking in 2D gels. Enough of each component is provided to process multiple protein samples. For researchers who have optimized an extraction protocol using one chaotropic extraction reagent, each kit reagent is also available as an individual product.

Features and Benefits:

- **Innovative detergent preparations** — Improved solubility allows for higher protein loads and greater visibility of low abundance proteins in 2D gels.
- Pre-mixed solubilization solutions

Ordering Information

| Description | Catalog No. |
|--|----------------|
| ProteoPrep® Total Extraction Sample Kit | PROTTOT |
| ProteoPrep® Universal Extraction Kit | PROTTWO |
| ProteoPrep® Membrane Extraction Kit | PROTMEM |
| ProteoPrep® Detergent Sample Kit | PROTDT |
| ProteoPrep® Reduction and Alkylation Kit | PROTRA |

Protein Extraction with Inhibitors

Featured Products

Protease Inhibitor Cocktails ▶

Prevent protein degradation by proteases during extraction and purification

Ensure the integrity of purified proteins by using protease inhibitor cocktails and highly specific protease inhibitors. During protein expression and isolation, endogenous proteases rapidly begin to degrade protein samples, reducing the quality and quantity of protein samples required for characterization and analysis. By using the right combination of protease inhibitors, you can protect your purified protein preparations from common proteases including serine proteases, metalloproteases, cysteine proteases, aminopeptidases, and aspartic proteases.

Ordering Information

| Application | Description | Catalog No. |
|--------------------------------------|---|--------------|
| SIGMAFAST™ Tablets | SIGMAFAST™ Protease Inhibitor Tablets, For General Use | S8820 |
| | SIGMAFAST™ Protease Inhibitor Cocktail Tablets, EDTA-Free, for use in purification of Histidine-tagged proteins | S8830 |
| For General Use | Protease Inhibitor Cocktail, for general use, lyophilized powder | P2714 |
| For Bacterial Extracts | Protease Inhibitor Cocktail, for use with bacterial cell extracts, lyophilized powder | P8465 |
| For Mammalian Cell & Tissue Extracts | Protease Inhibitor Cocktail, for use with mammalian cell and tissue extracts, DMSO solution | P8340 |
| For HIS-Tagged Proteins | Protease Inhibitor Cocktail, for use in purification of Histidine-tagged proteins, DMSO solution | P8849 |
| For Tissue Culture | Protease Inhibitor Cocktail, for use in tissue culture media, DMSO solution | P1860 |
| For Plant Extracts | Protease Inhibitor Cocktail, for plant cell and tissue extracts, DMSO solution | P9599 |
| For Fungal & Yeast Extracts | Protease Inhibitor Cocktail, for use with fungal and yeast extracts, DMSO solution | P8215 |

Roche cComplete™ Inhibitors

Roche offers a broad selection of protease inhibitors, as well as optimized lysis reagents, to ensure maximum yields of intact and functional proteins.

Don't spend valuable time and money repeating experiments in order to obtain sufficient yields of intact, functional proteins. Insist on Roche's high-quality protease inhibitors and lysis reagents to maximize success when isolating and purifying proteins.

Convenience

- Inhibit proteolytic activity in extracts from almost any tissue or cell type, including animals, plants, yeast, bacteria, and fungi.
- Drop a quick-dissolving tablet into your lysis buffer and eliminate the cumbersome job of weighing small amounts of different protease inhibitors on an analytical scale and dissolving the mix in DMSO.

Reliability

- Obtain stable, non-toxic protection in aqueous buffers.
- Consistently inhibit a multitude of protease classes, including serine proteases, cysteine proteases, and metalloproteases.

Enjoy Complete Protection

With eight powerful inhibitors, cComplete™ ULTRA Tablets reliably protect against a broad range of proteases for your most vital protein applications. For routine protein analyses, rely on the proven performance and convenience of classic cComplete™ Tablets from Roche. If your research requires inhibition of both proteases and phosphatases, combine cComplete™ ULTRA or cComplete™ Tablets with PhosSTOP™ Tablets for comprehensive protection.

Ordering Information

| Description | Qty | Catalog No. |
|---|----------------|--------------------|
| cComplete™ Tablets | 20 tablets | 11697498001 |
| | 3 x 20 tablets | 11836145001 |
| cComplete™ Tablets, EASYpack | 20 tablets | 4693116001 |
| cComplete™ Tablets, EDTA-free | 20 tablets | 11873580001 |
| | 3 x 20 tablets | 5056489001 |
| cComplete™ Tablets, EDTA-free, EASYpack | 20 tablets | 4693132001 |
| cComplete™ Tablets ULTRA, EDTA-free | 2 x 10 tablets | 5892953001 |
| | 6 x 10 tablets | 6538282001 |
| cComplete™ Tablets ULTRA | 2 x 10 tablets | 5892988001 |
| | 6 x 10 tablets | 6538304001 |
| cComplete™ Tablets, Mini | 25 tablets | 11836153001 |
| cComplete™ Tablets, Mini, EASYpack | 30 tablets | 4693124001 |
| cComplete™ Tablets, Mini, EDTA-free | 25 tablets | 11836170001 |
| cComplete™ Tablets, Mini, EDTA-free, EASYpack | 30 tablets | 4693159001 |
| cComplete™ Tablets ULTRA, mini, EASYpack | 30 tablets | 5892970001 |
| cComplete™ Tablets ULTRA, mini, EDTA-free, EASYpack | 30 tablets | 5892791001 |

Choose easy-to-use, versatile cOplete™ Protease Inhibitor Cocktail Tablets to obtain the protection you need, with convenience and reliability. Try cOplete™ Tablets today, and see how simple success can be.

cOplete™ ULTRA or cOplete™ Tablets selection guide

| | Performance | | | | Convenience | |
|-----------------------------------|------------------|--------------------|--------------------|------------------|--------------------------|-------------------------------|
| | Serine Proteases | Cysteine Proteases | Aspartic Proteases | Metalloproteases | Stability stock solution | Concentration stock solution* |
| cOplete™ ULTRA Tablets, EDTA-free | +++ | ++ | + | | ++ | 2x |
| cOplete™ ULTRA Tablets | +++ | ++ | + | + | ++ | 2x |
| cOplete™ Tablets, EDTA-free | ++ | + | | | +++ | 25x |
| cOplete™ Tablets | ++ | + | | + | +++ | 25x |

Phosphatase Inhibitor Cocktails

Prevent protein dephosphorylation for cell signaling studies

It is critical to preserve the phosphorylation state of proteins of interest during their extraction from cell and tissue lysates. To effect cell signaling, target proteins are phosphorylated by protein kinases that transfer a phosphate group to specific sites, typically at serine, threonine, or tyrosine residues. These phosphate groups can be removed by endogenous protein phosphatases,

restoring the protein to its original dephosphorylated state. Using phosphatase inhibitors can reveal the signaling status inside a cell at a specified timepoint. We offer a variety of Phosphatase Inhibitor cocktails and a PhosphoSafe™ Extraction Reagent that help protect phosphoproteins from different families of phosphatases.

Roche PhosSTOP™ Phosphatase Inhibitor Tablets

Achieve immediate, effective, and convenient inhibition of a broad spectrum of phosphatases across a wide range of sample materials (mammalian, plant, yeast, and bacteria) with non-toxic PhosSTOP™ Phosphatase

Inhibitor Cocktail Tablets. Cited in thousands of peer-reviewed papers, PhosSTOP™ Tablet is a proprietary blend of phosphatase inhibitors, formulated as a ready-to-use, quick-dissolving, water-soluble tablet.

Ordering Information

| Description | Recommended Application | Catalog No. |
|---|---|-------------|
| Phosphatase Inhibitor Cocktail Set I | Protection against alkaline phosphatases and Ser/Thr phosphatases such as PP1 and PP2A | 524624 |
| Phosphatase Inhibitor Cocktail Set II | Protection against acid and alkaline phosphatases and Protein Tyrosine Phosphatases (PTPs) | 524625 |
| Phosphatase Inhibitor Cocktail Set III | Protection against acid, alkaline and Ser/Thr phosphatases and Protein Tyrosine Phosphatases (PTPs) | 524627 |
| Phosphatase Inhibitor Cocktail Set IV | Protection against alkaline phosphatases and Ser/Thr phosphatases such as PP1 and PP2A | 524628 |
| PhosSTOP™ Phosphatase Inhibitor Tablets | Broad-spectrum blend of mammalian, plant, yeast, and bacterial phosphatase inhibitors | 4906845001 |

PROTEIN PURIFICATION AND DEPLETION

Affinity purification is based on the specific interaction of a target molecule with an immobilized ligand. We offer a wide range of tools for protein purification, including affinity magnetic beads, affinity agarose resins, the Amicon® Pro purification system and protease cleavage enzymes. To ensure that samples are enriched for protein(s) of interest, our depletion reagents eliminate common irrelevant, abundant proteins that may confound protein analysis. ●



- PureProteome™ magnetic beads are ideal for small volume affinity purification assays, such as immunoprecipitation and serum depletion or enrichment.
- Affinity agarose portfolio for larger volume applications, such as antibody purification and recombinant protein purification.
- Amicon® Pro purification system is ideal for small volume affinity purification assays followed by buffer exchange and/or concentration.
- Protease cleavage enzymes available in restriction grade or in kits for cleaving fusion proteins.



Affinity Purification with PureProteome™ Magnetic Beads ●

PureProteome™ Protein A and G Beads: Fast and easy immunoprecipitation

Traditional methods require hours of incubation time and harsh centrifugation to isolate sample. In contrast, PureProteome™ magnetic beads enhance binding equilibrium, enabling faster, gentler processing. The beads are easily resuspended for fast mixing and efficient interaction between the beads and protein.

PureProteome™ Protein A/G Mix Beads

Bind all mammalian immunoglobulin G (IgGs) efficiently using PureProteome™ Protein A/G mix magnetic beads, which provide a 50:50 blend of Protein A and Protein G.


Advantages of PureProteome™ Immunoprecipitation:

- **Be efficient with high capacity beads:** increased surface area allows for significantly greater binding capacity than other beads
- **Achieve high purity:** low non-specific binding of irrelevant proteins
- **Save time with fast sample processing:** enhanced binding equilibrium decreases incubation times by > 50%

PureProteome™ NHS and Carboxy FlexiBind beads

Customize your beads quickly and easily

Tailor your beads to match your application. Studying protein-protein interactions? Immobilizing enzymes, nucleic acids or small molecules? PureProteome™ NHS and Carboxy FlexiBind magnetic beads offer you flexibility in binding your target ligand. Customization of beads requires only that the target ligand has a free amine group.



PureProteome™ NHS FlexiBind Magnetic Beads (perfect for the first time user)

- **Fast:** Customize your own beads in < 60 min
- **Easy to Use:** Kit contains everything you need: beads, all buffers and Amicon® Ultra centrifugal filters for eliminating unreacted species
- **Robust:** Little experience or optimization required

PureProteome™ Carboxy FlexiBind Magnetic Beads (for the experienced user)


- **Flexible:** Choice of 0.3 µm, 1 µm or 2.5 µm COOH magnetic beads
- **Automation-Compatible:** Smaller beads have higher buoyancy properties while retaining strong magnetic capability

- **Flexibility:** Choose from a range of sizes and chemistries to fit your application
- **Cost Savings:** Less sample and reagent waste

PureProteome™ Beads

High speed immunoprecipitation with magnetic beads compared to agarose. In parallel indirect immunoprecipitations, PureProteome™ magnetic beads offered a 50% reduction in incubation time while yielding results equivalent to agarose beads.

Excellent yields.
Exceptional purity.
Faster protocol.



PureProteome™ Kappa and Lambda Ig Binder Beads

Immunoprecipitate all Human Antibodies (including IgA, IgD, IgE and IgM)

PureProteome™ Kappa Magnetic Beads bind to the kappa light chain constant region on human immunoglobulins with high specificity, and Lambda Magnetic Beads bind to the lambda light chain constant region. These novel magnetic beads are capable of capturing all immunoglobulin subtypes (IgG, IgA, IgD, IgE, and IgM) and provide a rapid, scalable, and reproducible means

to capture human antibody or antibody fragments containing kappa or lambda light chains — including F(ab) and F(ab')₂.

Depletion of all human immunoglobulins can be performed by mixing PureProteome™ Kappa and Lambda Magnetic Beads.

Relative Affinity

| | Protein A/G Mix | Protein A | Protein G | Kappa Ig Binder | Lambda Ig Binder | Kappa/Lambda Mix* |
|-------------------------|-----------------|-----------|-----------|-----------------|------------------|-------------------|
| Antibodies | | | | | | |
| Rabbit IgG | ● | ● | ● | | | |
| Mouse IgM | ● | ● | | | | |
| Mouse IgG ₃ | ● | ● | ● | | | |
| Mouse IgG _{2b} | ● | ● | ● | | | |
| Mouse IgG _{2a} | ● | ● | ● | | | |
| Mouse IgG ₁ | ● | ● | ● | | | |
| Human IgM | ● | ● | | ● | ● | ● |
| Human IgE | ● | ● | | ● | ● | ● |
| Human IgD | ● | ● | | ● | ● | ● |
| Human IgA | ● | ● | | ● | ● | ● |
| Human IgG ₄ | ● | ● | ● | ● | ● | ● |
| Human IgG ₃ | ● | | ● | ● | ● | ● |
| Human IgG ₂ | ● | ● | ● | ● | ● | ● |
| Human IgG ₁ | ● | ● | ● | ● | ● | ● |
| Rat IgM | ● | ● | | | | |
| Rat IgG _{2c} | ● | ● | ● | | | |
| Rat IgG _{2b} | ● | ● | ● | | | |
| Rat IgG _{2a} | ● | ● | ● | | | |
| Rat IgG ₁ | ● | ● | ● | | | |
| Rat IgG | ● | ● | ● | | | |


* PureProteome™ Kappa/Lambda mix is not a catalog item. Simply procure the Kappa and Lambda beads individually and mix at a 1:1 ratio.

| | A/G Mix | Protein A | Protein G | Kappa Ig Binder | Lambda Ig Binder | Kappa/Lambda Mix* |
|---------------------------|---------|-----------|-----------|-----------------|------------------|-------------------|
| Fragments | | | | | | |
| Human I | | | | | ● | ● |
| Human k | | | | ● | | ● |
| Human Fc | ● | ● | ● | | | |
| Human scFv | ● | ● | | ● | ● | ● |
| Human F(ab') ₂ | ● | ● | ● | ● | ● | ● |
| Human F(ab) | ● | ● | ● | ● | ● | ● |

Key code for relative affinity of protein A and G; PureProteome™ Kappa and Lambda magnetic beads for respective antibodies:

- Strong Affinity
- Moderate/Slight Affinity
- Requires Evaluation

PureProteome™ Kappa or Lambda light chain ligands bind to the constant region of the antibody light chain, and will not bind the scFv.



Ordering Information

| Application | Description | Catalog No. |
|---|--|--------------|
| IP, Antibody Purification, F(ab) Purification | PureProteome™ Protein A Magnetic Beads | LSKMAGA10 |
| | PureProteome™ Protein G Magnetic Beads | LSKMAGG10 |
| | PureProteome™ Protein A/G Mix Magnetic Beads | LSKMAGAG10 |
| | PureProteome™ Kappa Ig-Binder Magnetic Beads* | LSKMAGKP02 |
| | PureProteome™ Lambda Ig-Binder Magnetic Beads* | LSKMAGLM02 |
| Biotinylated Molecule Purification | PureProteome™ Streptavidin Magnetic Beads | LSKMAGT10 |
| His•Tag® Tagged Protein Purification | PureProteome™ Nickel Magnetic Beads | LSKMAGH10 |
| Custom Labelled (Flexibility to Bind Ligand of Choice) | PureProteome™ NHS FlexiBind Magnetic Beads | LSKMAGN04 |
| | PureProteome™ Carboxy FlexiBind Magnetic Beads** | LSKMAG1CBX10 |
| Magnetic Stands | PureProteome™ Magnetic Stand, 8-well | LSKMAGS08 |
| | PureProteome™ Magnetic Stand, 15 mL | LSKMAGS15 |

*Human only.

**Available in 0.3, 1.0 and 2.5 µm.

Agarose Based Affinity Purification ▶

Agarose resins are the preferred approach for large purifications and a convenient option when scaling up will be needed. We offer a complete portfolio of agarose resins and kits for antibody purification, immunoprecipitation, and purification of tagged proteins.

Ordering Information

| Description | Size | Catalog No. |
|-------------------------------|--------|-------------|
| Protein A Agarose | 1.5 mL | IP02-1.5ML |
| | 10 mL | 16-125 |
| Protein A Agarose Fast Flow | 10 mL | 16-156 |
| Protein G Agarose | 1.5 mL | IP04-1.5ML |
| | 10 mL | 16-266 |
| Protein A + Protein G Agarose | 1.5 mL | IP05-1.5ML |
| | 10 mL | IP10-10ML |

Affinity Purification with Recombinant Fusion Tags ▶

Recombinant protein purification by tag-specific affinity chromatography is a proven technology that results in highly specific recognition and purification of recombinant proteins. Our broad line of purification and detection tools includes the HIS-Select®, GST, HA, and FLAG® reagents and other technologies.

FLAG® System ▶

The FLAG® Expression System is a tested method for expression, purification, and detection of recombinant fusion proteins. The FLAG® and 3x FLAG®

Antibody Purification and Immunoprecipitation ▶

Protein A and Protein G are proteins of microbial origin that bind specifically to mammalian immunoglobulins. When coupled to agarose, they provide an efficient tool for purification and immunoprecipitation of antibodies. Immunoglobulins of various species interact differently with the two proteins. Agarose that combines Protein A and Protein G provides the binding characteristics of both in a single reagent.

systems are useful in western blotting, immunocytochemistry, immunoprecipitation, flow cytometry, protein purification, and in the study of protein-protein interactions. These small hydrophilic tags significantly improve detection and purification of recombinant fusion proteins when used with our highly specific and sensitive anti-FLAG® antibodies. Sensitivity can be enhanced up to 200-fold using the 3x FLAG® epitope.

- Sequence is highly charged and useful for sensitive detection
- Sensitivity can be enhanced using the 3x FLAG® epitope
- Enhances the study of low-abundance proteins and the optimization of difficult protein expression projects

Ordering Information

| Description | Pack Size/Quantity | Catalog No. |
|---|--|-------------|
| Purification | | |
| FLAG® Affinity Gels | | |
| ANTI-FLAG® M1 Agarose Affinity Gel | 1 mL, 5 mL, 10 mL, 25 mL | A4596 |
| ANTI-FLAG® M2 Affinity Gel, Purified Immunoglobulin, Buffered Aqueous glycerol solution | 1 mL, 5 mL, 10 mL, 25 mL, 2 x 25 mL, 4 x 25 mL | A2220 |
| EZview™ Red ANTI-FLAG® M2 Affinity Gel | 500 µL, 1 mL, 5 X 1 mL | F2426 |
| FLAG® Magnetic Beads | | |
| ANTI-FLAG® M2 Magnetic Beads | 1 mL, 5 mL | M8823 |
| FLAG® 96-well Format | | |
| FLAG® M Purification Kit, For Mammalian expression systems | 1 Kit | CELLMM2 |
| ANTI-FLAG® High Sensitivity, M2 coated 96-well plates, 96-well, clear, polystyrene, flat bottom plate | 1 Each, 5 Each, 100 Each | P2983 |
| FLAG® HA Affinity Purification Kits | | |
| FLAG® HA Tandem Affinity Purification Kit | 5 Reactions | TP0010 |
| FLAG® Peptides | | |
| FLAG® Peptide, lyophilized powder | 4 mg, 25 mg | F3290 |
| 3X FLAG® Peptide, lyophilized powder | 1 mg, 4 mg, 25 mg | F4799 |

HIS-Select® System ►

Our HIS-Select® products purify histidine-tagged proteins quickly and with high selectivity made possible by a patented HIS-Select® metal chelate linker which is hydrophilic and non-charged. Because the HIS-select® linkage chemistry is uncharged, non-specific binding of unwanted proteins is dramatically reduced. In addition, the novel tetradentate chelator used in the HIS-Select® products reduces nickel leaching from the affinity gel, affording higher purity and binding capacity for histidine-tagged proteins. The need for secondary purification of HIS-Select® tagged proteins is eliminated due to a single-step purification procedure.

Features and Benefits

- One-step purification
- High selectivity for enhanced purity of target proteins
- Non-charged, hydrophilic linkage reduces non-specific binding
- Pure tetradentate chelate for minimal metal leaching

Ordering Information


| Description | Pack Size/Quantity | Catalog No. |
|--|------------------------------------|-------------|
| HIS-Select® Affinity Gels | | |
| HIS-Select® Nickel Affinity Gel | 1 mL, 5 mL, 25 mL, 100 mL, 500 mL | P6611 |
| HIS-Select® HF Nickel Affinity Gel, High Flow | 1 mL, 10 mL, 25 mL, 100 mL, 500 mL | H0537 |
| HIS-Select® Cobalt Affinity Gel | 5 mL, 25 mL, 100 mL | H8162 |
| IMAC-Select Affinity Gel | 5 mL, 25 mL, 100 mL | I1408 |
| EZview™ Red HIS-Select® HC Nickel Affinity Gel | 1 mL, 5 X 1 mL | E3528 |
| HIS-Select® Convenient Formats | | |
| HIS-Select® Spin Columns | 10 Each, 50 Each | H7787 |
| HIS-Select® 96-well Format | | |
| HIS-Select® Filter Plate | 1 Each, 5 X 1 Each | H0413 |
| HIS-Select® Buffers | | |
| HIS-Select® Wash Buffer | 500 mL, 1 L | H5288 |
| HIS-Select® Elution Buffer | 250 mL, 500 mL | H5413 |

His•Tag® Purification

Ni-NTA His•Bind® Resin has a binding capacity of over 10 mg of His-Tagged fusion protein per mL resin.

The agarose matrix on the Ni-NTA His•Bind® Superflow™ Resin is structured with more crosslinking for enhanced bead rigidity, for exceptional compatibility with FPLC.

Our IDA His•Bind® resins are offered uncharged to allow flexibility of choice in the metal ion (Nickel, Cobalt, Zinc, Iron, Copper, etc.). IDA supports can be recycled many times with no loss in performance.



| Lane | Sample |
|------|-----------------------------|
| 1 | Crude Extract |
| 2 | Markers |
| 3 | Ni-NTA Competitor Q Elution |
| 4 | Ni-NTA Competitor Q Strip |
| 5 | Ni-NTA Competitor G Elution |
| 6 | Ni-NTA Competitor G Strip |
| 7 | Ni-NTA His•Bind® Elution |
| 8 | Ni-NTA His•Bind® Strip |

Ni-NTA His•Bind® performance vs. equivalent competitor resins Vector pET-28b (+) was used to express a His-Tag fusion protein of 119KDa in E. coli BL21 (DE3) cells, induced culture was processed with BugBuster® Master Mix, and protein extract was divided evenly to proceed to the His-Tag purification using Ni-NTA His•Bind®, Ni-NTA Competitor Q and Ni-NTA Competitor G resins. Ni-NTA His•Bind® resins show higher binding capacity and a better purification.

Ordering Information

| Application | Description | Catalog No. |
|--|--|-------------|
| Ni-NTA His•Bind® Resin | | |
| Small to medium scale | Ni-NTA His•Bind® Resin | 70666 |
| Gravity flow column | BugBuster® Ni-NTA His•Bind® Purification Kit | 70751 |
| Recommended for eukaryotic extracts | Ni-NTA Buffer Kit | 70899 |
| Ni-NTA His•Bind® Superflow™ Resin | | |
| Small to production scale | Ni-NTA His•Bind® Superflow™ Resin | 70691 |
| FPLC or gravity flow column | Ni-NTA Buffer Kit | 70899 |
| Uncharged IDA His•Bind® Resin | | |
| Uncharged (metal flexibility) | IDA His•Bind® Resin | 69670 |
| Reusability | His•Bind® Buffer Kit | 69755 |
| Small to medium scale | His•Bind® Purification Kit | 70239 |
| Gravity flow column or batch mode | BugBuster® His•Bind® Purification Kit | 70793 |

GE HisTrap™ Columns

Prepacked HisTrap™ columns from GE Healthcare deliver maximum convenience for greater flexibility and reduced hands-on operation.

- Simple, high resolution purification of histidine-tagged proteins
- Minimized risk of deactivation of target proteins due to broad compatibility with a wide range of reducing agents, detergents, and other additives
- HisTrap™ Fast Flow Column delivers fast flow rate purification and easy scale-up

Ordering Information

| Description | Qty | Catalog No. |
|--|------------|--------------|
| Prepacked columns and kit | | |
| HisTrap™ High Performance Column | 5 x 1 mL | GE17-5247-01 |
| HisTrap™ High Performance Column | 1 x 5 mL | GE17-5248-01 |
| HisTrap™ High Performance Column | 5 x 5 mL | GE17-5248-02 |
| HisTrap™ High Performance Column | 100 x 5 mL | GE17-5248-05 |
| HisTrap™ High Performance Kit | 1 Kit | GE17-5249-01 |
| For high flow rate purification, scale-up and manual purification | | |
| HisTrap™ Fast Flow Column | 5 x 1 mL | GE17-5319-01 |
| HisTrap™ Fast Flow Column | 5 x 5 mL | GE17-5255-01 |
| HisPrep™ Fast Flow 16/10 Column | 1 x 20 mL | GE28-9365-51 |

Hemagglutinin (HA) Purification & Detection ▶


Exceptional sensitivity and specificity for detection of Hemagglutinin (HA) — tagged proteins. The HA peptide is suitable for use in immunoblotting or to elute HA-tagged fusion proteins.

Ordering Information

| Description | Pack Size/Qty | Catalog No. |
|---|------------------------|-------------|
| Influenza Hemagglutinin (HA) Peptide, ≥ 97% (HPLC) | .5 mg, 1 mg | I2149 |
| Monoclonal Anti-HA antibody produced in mouse purified immunoglobulin, clone HA-7 | 200 µL | H3663 |
| Monoclonal Anti-HA–Agarose antibody produced in mouse clone HA-7, purified immunoglobulin | 1 mL, 5 x 1 mL | A2095 |
| Anti-HA Immunoprecipitation | 1 Kit | IP0010 |
| Monoclonal Anti-HA–Peroxidase antibody produced in mouse, clone HA-7, purified immunoglobulin, lyophilized powder | 1 Vial | H6533 |
| EZview™ Red Anti-HA Affinity Gel | 500 µL, 1 mL, 5 x 1 mL | E6779 |

GST•Tag™ Purification

The GST fusion system is based on the widely recognized affinity of glutathione-S-transferase (GST) fusion proteins for immobilized glutathione. Our GST Resin utilizes an 11-atom spacer arm to covalently attach reduced glutathione to the solid support via a sulfide linkage. The resin can be reused several times without loss of capacity, and the high degree of substitution of glutathione ensures exceptional binding capacity.



| Lane | Sample |
|------|------------------------------------|
| M | PerfectProtein™ markers 15–150 kDa |
| 1 | BugBuster® extract |
| 2 | Flow-through |
| 3 | Eluate |

GST•TAG™ Purification

GST•Bind™ purification. A crude extract containing unfused GST was applied to a 2 mL GST•Bind™ Resin column. Total protein yield after purification was 8 mg/mL resin.

S•Tag™ Purification

The S•Tag™ fusion protein is a short 15-aa sequence that specifically binds with high affinity the 104-aa S-Protein (KD = 10⁻⁹ M, 1000 times stronger than the interaction between nickel and His•Tag® fusion protein). Fusion proteins can be easily purified by cleavage with site-specific proteases or in acidic buffers.



| Lane | Sample |
|------|--|
| M | PerfectProtein™ markers 15–150 kDa |
| 1 | Crude extract |
| 2 | Flow-through |
| 3 | Wash 1 |
| 4 | Wash 2 |
| 5 | Eluate + Biotinylated Thrombin |
| 6 | Eluate after Biotinylated Thrombin removal |

S•Tag™ affinity purification

S•Tag™ β-gal expressed from a pET construct was purified from a crude soluble fraction using S-protein Agarose under native conditions. Elution of the target protein from the agarose was performed by digestion with Biotinylated Thrombin, which was subsequently removed with Streptavidin Agarose. The fractions are indicated.

Strep•Tag® II Purification

The Strep•Tag® fusion protein II is an 8 amino acid sequence that binds to the biotin pocket of Streptavidin with 100 times higher binding capacity.

T7•Tag® Purification

Purification is antibody-based. Covalently coupled to agarose beads, the T7•Tag® monoclonal antibody captures the T7•Tag® epitope — a sequence of 11 amino acids.

Streptavidin Agarose

Cross-linked agarose is covalently coupled with pure streptavidin under controlled conditions. The stable linkage to the resin minimizes leaching of the streptavidin while maintaining full binding activity. The matrix is suitable for use in column and batch formats for any application that requires high biotin binding capacity and low non-specific binding, and is ideal for affinity purification of biotinylated proteins or pull down experiments of biotinylated DNA/RNA probes. The resin has no detectable protease, DNase, or RNase.

Ordering Information

| Description | Catalog No. | |
|---------------------------------------|-------------|---------|
| GST•Tag™ Purification | | |
| GST•Bind™ Resin | 70541 | |
| GST•Bind™ Buffer Kit | 70534 | |
| BugBuster® GST•Bind™ Purification Kit | 70794 | |
| S-Tag Purification | | |
| S-protein Agarose | 69704 | |
| S•Tag™ Thrombin Purification Kit | 69232 | |
| S•Tag™ rEK Purification Kit | 69065 | |
| Strep•Tag® II Purification | | |
| Strep-Tactin® Superflow Agarose | 71592 | |
| Strep-Tactin® Buffer Kit | 71613 | |
| Strep-Tactin® SpinPrep Kit | 71608 | |
| D-Desthiobiotin | 71610 | |
| T7•Tag® Purification | | |
| T7•Tag® Affinity Purification Kit | 69025 | |
| T7•Tag® Antibody Agarose | 69026 | |
| Description | | |
| Streptavidin Agarose | 5 mL | 69023-3 |
| | 10 mL | 16-126 |

Protein Depletion

Seppro® Protein Depletion Technology

Overcome protein sample complexity — Separate with Seppro® Depletion Technology

Seppro® Depletion Technology enables removal of highly abundant proteins that may mask target protein detection from a variety of biological samples using the affinity of avian polyclonal IgY antibodies.

The Seppro® platform, incorporating Supermix technology, represents the most complete human protein depletion

system available, removing 14 of the most abundant proteins from human serum or plasma, as well as other high and medium abundance proteins. Additional products are available for the depletion of mouse and rat samples, as well as the industry's only depletion system for the removal of Rubisco from plant samples.

Ordering Information

| Description | Capacity | Uses | Proteins Depleted | Catalog No. |
|------------------------|------------------------------|------|--|-------------|
| Human | | | | |
| IgY14 Spin Columns | 15–20 µL | 200 | 14 most abundant proteins | SEP010 |
| IgY14 LC2 | 40–50 µL | 100 | | SEP020 |
| IgY14 LC5 | 100 µL | 100 | | SEP030 |
| IgY14 LC10 | 200–250 µL | 100 | | SEP040 |
| HT IgY14 96 well plate | 1.5–2.0 µL per well | 10 | | S2453 |
| Human Supermix LC2 | Flow through from IgY14 LC5 | 100 | Moderately abundant proteins, resulting in 99% total protein removal | SEP050 |
| Human Supermix LC5 | Flow through from IgY14 LC10 | 100 | | SEP060 |
| Rat | | | | |
| Rat Spin Columns | 15–20 µL | 200 | 7 most abundant proteins | SEP130 |
| Rat LC 10 | 200–250 µL | 100 | | SEP120 |
| Mouse | | | | |
| Mouse Spin Columns | 15–20 µL | 200 | 7 most abundant proteins | SEP110 |
| Mouse LC10 | 200–250 µL | 100 | | SEP090 |
| Mouse Supermix LC5 | Flow-through from mouse LC10 | 100 | Further partitions complex mouse plasma/serum samples | SEP100 |
| Plant | | | | |
| RuBisCO Spin Columns | 15–20 µL | 200 | RuBisCO (Ribulose-1,5- bisphosphate carboxylase/oxygenase) | SEP070 |
| RuBisCO LC2 | 40–50 µL | 100 | | SEP080 |

ProteoExtract® Agarose Columns

Human serum and plasma samples are rich sources of proteomic information, reflecting processes regulating normal or diseased states. Today's ultra-sensitive analytical methods, such as two-dimensional (2D) gel electrophoresis and mass spectrometry, can detect minute changes in expression profiles — but ultrasensitive approaches typically require the removal of highly abundant proteins (HAP) and moderately abundant proteins (MAP).

We offer a range of kits and resins for depleting high-abundance proteins (HAP) from serum or plasma samples. Choose from the PureProteome™ magnetic bead kits and resins or the ProteoExtract® agarose columns. First, identify the species of your serum/plasma source — the following tables summarize the different solutions for your needs.

Ordering Information

| Description | Format | Species | Proteins Depleted | Catalog No. |
|--|-----------------|---|-----------------------|---------------|
| ProteoExtract® Albumin/IgG Depletion Kit | Agarose columns | Human, rabbit, rat, mouse, pig and bovine | Albumin and IgG > 80% | 122642 |
| ProteoExtract® Albumin Depletion Kit | | | Albumin > 80% | 122640 |

| Application | Description | Catalog No. |
|----------------------|---|--------------------|
| Depletion/Enrichment | PureProteome™ Albumin Magnetic Beads | LSKMAGL10 |
| | PureProteome™ Albumin/IgG Depletion Kit | LSKMAGD12 |
| | PureProteome™ Human Albumin/Immunoglobulin Depletion Kit* | LSKMAGHDKIT |

*Human only.

ProteoPrep® Immunoaffinity Albumin & IgG Depletion Kit

The ProteoPrep® Immunoaffinity medium in prepacked spin columns is a mixture of two-beaded media containing recombinantly expressed, small single-chain antibody ligands, providing low non-specific binding and high capacity. This kit is also effective in depleting albumin and IgG from mouse and guinea pig serum.

The ProteoPrep® Blue Albumin & IgG Depletion Kit is designed to specifically remove albumin and IgG from 25 samples of human serum (25 µL to 100 µL) in preparation for two-dimensional electrophoresis (2DE). Albumin-depleted serum samples generated by this kit are in urea rather than salt-based buffers with little dilution, eliminating the need to precipitate the sample prior to performing 2DE.

Ordering Information

| Description | Format | Proteins Depleted | Catalog No. |
|--|-------------|--------------------------|-------------|
| ProteoPrep® Immunoaffinity Albumin & IgG Depletion Kit | Spin Column | Albumin > 95%, IgG > 85% | PROTIA |
| ProteoPrep® Blue Albumin & IgG Depletion Kit | | Albumin > 85%, IgG > 70% | PROTBA |


Amicon® Pro Purification System

Purify biologically active proteins with gentle, all-in-one recovery.

Biologically active proteins yield meaningful data. When you start with consistent yields of active, native-folded protein, you're giving your experiment the best chance to succeed. If your current protein preps involve juggling columns, dialyzers and multiple transfer steps, you could be introducing variability into your data. For your next protein preparation, choose the simple, gentle method that tackles even the most labile and poorly expressed proteins — the Amicon® Pro purification system. When your proteins behave, your research will flourish.



WORKING WITH PROTEINS?



A simple, flexible tool for the basic researcher.

Whether you're performing affinity purification from a precious sample, labeling antibodies, depleting abundant proteins from serum samples or removing salts from a chromatography sample, the Amicon® Pro system is your sample preparation partner. The modular design not only allows flexibility in application, but also offers unprecedented simplicity in protein sample preparation.

Examples:

- Turn your crude lysate into a purified, concentrated protein ready for your downstream assay in as few as four spins.
- Perform a 99% buffer exchange using a patent-pending, continuous, gentle process in one spin.


Don't lose protein in multiple devices.

Maximize your protein recovery with the Amicon® Pro System. ●

Traditional protein purification can be a long process with multiple steps and devices, which can often result in protein degradation and loss along the way. By using the Amicon® Pro Purification System, you can avoid the risks involved with sample transfer while reducing hands-on time.

Whether you need to affinity purify, concentrate, dialyze, or any combination of the three, the Amicon® Pro Purification System will save time and improve your protein recovery. It can help you perform multiple protein preparations in parallel, improving prep-to-prep reproducibility and enabling head-to-head comparison of expression constructs.

Amicon® Pro system unique design features and workflow benefits



"If I was doing things the old way, I would be six months — if not a year — behind where I am right now with my project."

-Jason Lehmann, Amicon® Pro user, University of California in San Diego

Ordering Information

To choose the appropriate Amicon® Pro device, determine the nominal molecular weight cut-off (NMWCO) for your protein of interest and your desired affinity purification scheme. For convenience and ease of use, the Amicon® Pro purification kits contain devices, reagents and buffers optimized for twelve reactions. These kits are ideal for affinity purification of tagged recombinant proteins, antibody purification and depletion.

| Amicon® Pro Purification Kits 12/pk Includes reagent kit (resin and buffers) | Reagent Kit Only | NMWCO | | | | |
|---|---------------------|-----------|-----------|-----------|-----------|-----------|
| | | 3,000 | 10,000 | 30,000 | 50,000 | 100,000 |
| Amicon® Pro Affinity Concentration Kit Ni-NTA | ACR5000NT | ACK5003NT | ACK5010NT | ACK5030NT | ACK5050NT | ACK5100NT |
| Amicon® Pro Affinity Concentration Kit Protein A | ACR5000PA | ACK5003PA | ACK5010PA | ACK5030PA | ACK5050PA | ACK5100PA |
| Amicon® Pro Affinity Concentration Kit Protein G | ACR5000PG | ACK5003PG | ACK5010PG | ACK5030PG | ACK5050PG | ACK5100PG |
| Amicon® Pro Affinity Concentration Kit GST | ACR5000GS | ACK5003GS | ACK5010GS | ACK5030GS | ACK5050GS | ACK5100GS |

| Amicon® Pro purification system — No Reagents Included | NMWCO | | | | |
|--|-----------|-----------|-----------|-----------|-----------|
| | 3,000 | 10,000 | 30,000 | 50,000 | 100,000 |
| Amicon® Pro Purification System Trial Pack 2/pk | ACS500302 | ACS501002 | ACS503002 | ACS505002 | ACS510002 |
| Amicon® Pro Purification System 12/pk | ACS500312 | ACS501012 | ACS503012 | ACS505012 | ACS510012 |
| Amicon® Pro Purification System 24/pk | ACS500324 | ACS501024 | ACS503024 | ACS505024 | ACS510024 |
| Amicon® Pro Purification System 24/pk without Amicon® Ultra 0.5 mL filter: | ACS500024 | | | | |

Protein Purification with Protease Cleavage Enzymes


Featured Products

Restriction and Biotinylated Grade Thrombin ●

Highly efficient, specific cleavage of fusion proteins

Restriction Grade Thrombin is qualified to specifically cleave target proteins containing the recognition sequence LeuValProArg↓GlySer. The preparation is functionally tested for activity with fusion proteins and is free of detectable contaminating proteases. Thrombin is supplied with 10X Thrombin Cleavage Buffer and a Cleavage Control Protein.

Biotinylated Thrombin is identical in activity to Restriction Grade Thrombin, but has covalently attached biotin for easy removal of the enzyme from cleavage reactions using immobilized streptavidin. Our Thrombin Cleavage Capture Kit includes not only biotinylated thrombin and immobilized streptavidin, but also all required buffers and filters for complete, convenient recovery of cleaved protein.



Biotinylated Thrombin cleavage. The indicated amounts of Biotinylated Thrombin were used to cleave 2 μ g of Cleavage Control Protein in an overnight digestion. Samples were analyzed by SDS-PAGE (4–20% gradient gel) followed by staining with Coomassie® blue stain. The 0.0045 unit lane represents a 2.25-fold over-digestion.

HRV 3C Protease


Highly efficient, specific cleavage of fusion proteins

Recombinant type 14 3C protease from human rhinovirus (HRV 3C) is a highly purified, recombinant 6XHis-tagged enzyme, which recognizes the cleavage site LeuGluValLeuPheGln↓GlyPro.

The small, 22 kDa size of the protease, with optimal activity at 4 °C, high specificity, and His-tag fusion make HRV 3C protease an ideal choice for rapid removal of fusion tags.

| Lane | Sample |
|------|---|
| M | PerfectProtein Markers, 10–225 kDa |
| 1 | 3 µg purified Nus•Tag™ enolase fusion protein |
| 2 | 3 µg Nus•Tag™ enolase fusion protein with 30 min HRV3C protease reaction |
| 3 | 3 µg Nus•Tag™ enolase fusion protein with 30 min competitor's protease reaction |
| 4 | 3 µg Nus•Tag™ enolase fusion protein with 60 min HRV3C protease reaction |
| 5 | 3 µg Nus•Tag™ enolase fusion protein with 60 min competitor's protease reaction |

HRV 3C Protease cleaves fusion proteins more efficiently compared to cleavage with a competitor's protease. Using a 1:100 (w/w) ratio of protease:target protein, 500 µg of purified Nus•Tag™ enolase fusion protein was incubated in parallel 500 µL reactions at 4 °C. The reactions were quenched by adding equal volume 4X SDS Sample Buffer and then immediately placing the samples into a water bath at 75 °C for 5 min.



Ordering Information

| Description | Catalog No. |
|-----------------------------------|-------------|
| Restriction-Grade Thrombin | 69671 |
| Biotinylated Thrombin | 69672 |
| Thrombin Cleavage Capture Kit | 69022 |
| Restriction Grade Factor Xa | 69036 |
| Factor Xa Cleavage Capture Kit | 69037 |
| Recombinant Enterokinase | 69066 |
| Enterokinase Cleavage Capture Kit | 69067 |
| HRV 3C Protease | 71493 |
| Tag•off™ High Activity rEK | 71537 |
| Tag•off™ rEK Cleavage Capture Kit | 71540 |

PROTEIN BUFFER OPTIMIZATION AND SAMPLE CONCENTRATION

When downstream quality matters, make sure your upstream tools are the best. The last steps of preparing a protein sample for downstream analyses, such as activity assays or structural studies, involve ensuring that the protein is in its native, soluble form, dissolved in the buffer of choice, and at an appropriate concentration. With our tools for protein buffer optimization and sample concentration, obtain publication-quality data from every last microgram of protein.

Protein Buffer Exchange, Sample Desalting, and Dialysis ●

Each protein preparation is unique. Give it the special treatment it deserves with a perfectly designed device for dialyzing and buffer exchange. Select between fast and gentle diafiltration using the Amicon® Pro System or dialysis using D-Tube™ Dialyzers.

| Sample Needs | Amicon® Pro System | Amicon® Ultra Filter | D-Tube™ Dialyzer |
|--|---|---|--------------------------------|
| Faster optimization | ~20 minutes | < 1 hour | 5 hours |
| Sensitive samples which may precipitate at higher concentrations | + | - | + |
| Post-dialysis concentration | + | + | - |
| Limited amounts of exchange solvent | + | + | - |
| Temperature sensitive | Minimal effect of cold temperature on speed | Minimal effect of cold temperature on speed | Cold temperature reduces speed |

Novel engineering provides unmatched buffer exchange. ●

The Amicon® Pro device is the first of its kind to offer dynamic, continuous buffer exchange by diafiltration.


How does it work? The secret is in the design of the Amicon® Pro exchange device and tip. The lower portion of the exchange device is designed to exactly match the contours of the Amicon® Ultra-0.5 mL filter. The tip is tapered to maximize the external-to-internal volume ratio, ensuring that fresh buffer is slowly but consistently metered in, mixed with sample, and forced across the membrane and out. This delivers a continuous, controlled flow during desalting and buffer exchange, without multiple dilute-and-concentrate centrifugation steps. The results are the gentle recovery of greater than 95% of purified protein.

Fast: single spin

Gentle: unique design provides continuous diafiltration

Less Buffer: only 1.5 mL buffer required (given 0.5 mL initial sample)





The uniquely designed interface between the exchange tube tip and the Amicon® Ultra device enables greater than 99% buffer exchange in a single spin. Buffer exchange, as shown in this diagram, was measured by the replacement of a low-molecular weight dye (yellow) with clear buffer (black arrows); while a high-molecular weight dye (bright blue) was retained inside the Amicon® Ultra device.


The gentleness of dialysis with the efficiency of diafiltration.

| | Dialysis cassette + concentrator | 0.5 mL diafiltration device (3 spin) | Amicon® Pro purification system |
|---------------------------------------|----------------------------------|--------------------------------------|---------------------------------|
| Process time | 16 hours | 50 min. | 20 min. |
| Recovery | 51 % | > 95% | > 95 % |
| Specific activity (signal/µg GST-LPP) | 0.195 | 0.17 | 0.199 |

Gentler buffer exchange = greater activity. Eluted Samples of GST-lambda protein phosphatase (LPP) buffer exchanged and concentrated using Amicon® Pro device showed greater specific activity and percentage recovery than when prepared with a dialysis cassette (plus concentrator) or 0.5 mL diafiltration spin column.

One hour antibody labeling with the Amicon® Pro system. ●

The unique design of the exchange tip enables single spin diafiltration.



Generate FITC-labeled antibody in one hour. What's faster than labeling antibodies using other purification methods, and more economical than purchasing pre-labeled antibodies? Using Amicon® Pro purification systems for antibody labeling.

| Step | Dialysis-based buffer exchange pre/post labeling | Amicon® Pro purification system |
|---------------------------------------|--|---------------------------------|
| Buffer exchange | Overnight | 15 min |
| FITC labeling | 3 h | 30 min |
| Free FITC removal and buffer exchange | Overnight | 15 min |
| Total time | 3 days | 1 h |
| Antibody recovery | 39% | 72% |

Ordering Information

To choose the appropriate Amicon® Pro device, determine the nominal molecular weight cut-off (NMWCO) for your protein of interest and your desired affinity purification scheme.

| Amicon® Pro purification system — No Reagents Included | NMWCO | | | | |
|---|-----------|-----------|-----------|-----------|-----------|
| | 3,000 | 10,000 | 30,000 | 50,000 | 100,000 |
| Amicon® Pro Purification System 12/pk | ACS500312 | ACS501012 | ACS503012 | ACS505012 | ACS510012 |
| Amicon® Pro Purification System 24/pk | ACS500324 | ACS501024 | ACS503024 | ACS505024 | ACS510024 |
| Amicon® Pro Purification System 24/pk without Amicon® Ultra 0.5 mL filter | ACS500024 | | | | |

Featured Products

D-Tube™ Dialyzers

Fast and easy dialysis

Gently dialyze intractable or sensitive samples and prevent them from precipitation or over-concentration. Providing maximum efficiency, D-Tubes™ dialyzers are designed with a double membrane to spread the sample over a large surface area enabling complete dialysis in just two to five hours.



D-Tube™ Dialyzer Advantages:

> 89% Sample Recovery

- Low binding membrane and housing enhance sample recovery

Reliable and Easy to Use

- Secure design prevents sample loss due to leaks — no knots or clamps to loosen and leak
- Easy to open and close with a screw cap
- Rigid frame permits smooth sample withdrawal of submilliliter volumes — removing every last drop is easy

Convenient Sample Loading

- No need to use a syringe to load or remove samples. Simply load your sample with standard pipette tip
- Floating racks fit most standard beakers to hold devices in exchange buffer
- D-Tubes™ dialyzers can also be used to electroelute samples from agarose or acrylamide

Ordering Information

| | | Product | D-Tube™ Dialyzer Mini | D-Tube™ Dialyzer Midi | D-Tube™ Dialyzer Maxi | D-Tube™ Dialyzer Mega | D-Tube™ Dialyzer Mega |
|-----------------------------------|--------------------------|-------------------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| Proteins/DNA/RNA/Oligonucleotides | Molecular Weight Cut-off | Maximum initial sample volume | 10 to 250 µL | 50 to 800 µL | 100 µL to 3 mL | 3 to 10 mL | 10 to 15 mL |
| MW | MWCO (Da) | Qty/pk | | | | | |
| MW < 7 k | 3,500 | 10 | | 71506-3 | 71508-3 | 71739-3 | 71742-3 |
| | | 50 | | | | 71739-4 | 71742-4 |
| 7 < MW < 24 k | 7,000 | 10 | 71504-3 | 71507-3 | 71509-3 | 71740-3 | 71743-3 |
| | | 50 | | | | 71740-4 | 71743-4 |
| 24 k < MW | 13,000 | 1 plate of 96 | 71712-3 | | | | |
| | | 10 | 71505-3 | | 71510-3 | | |
| | | 50 | | | | | |
| | | 1 plate of 96 | 71713-3 | | | | |
| | Floating Rack | Product (Qty/pk) | Mini (10) | Midi (10) | Maxi (10) | Mega (10) | Mega (10) |
| | | | 71512-3 | 71513-3 | 71514-3 | 71748-3 | 71748-3 |

Centrifugal Concentration Devices

Fast and Easy Diafiltration with Amicon® Ultra Centrifugal Filters

Change buffers by gradually adding new solvent during simultaneous ultrafiltration

Because some macromolecules can lose activity or proper structure upon extreme changes of buffer conditions, use diafiltration, which involves removing microsolute by adding solvent to the sample being filtered at the same time that ultrafiltration is being applied.

Advantages of Amicon® Ultra Centrifugal Filters diafiltration:

- Fast — buffer exchange in as few as two spins
- Efficient — requires minimal volume of exchange buffer, easily contained in reservoir
- Easy to use — simply load your sample with standard pipette tip
- Enables simultaneous concentrating and desalting



Featured Products

Amicon® Ultra Centrifugal Filters

Fast and Easy Protein Concentration

Amicon® Ultra Centrifugal filters provide fast sample processing and promote high sample recoveries, even in dilute samples, through ultrafiltration. The unique features of the Amicon® Ultra centrifugal filters give you the fastest, most efficient concentration for sensitive downstream applications.

Amicon® Ultra Centrifugal Filter Advantages:

Maximize Concentration with Highest Protein Recovery – True Engineered Dead Stop

- Avoids spinning to dryness
- Provides a predictable concentration factor
- No need to calibrate for several samples to run in parallel

Reverse Spin Recovery

- Reverse spin devices enable you to maximize protein recovery, especially with small dilute samples, without introducing pipetting errors
- Low binding membrane and polypropylene housing for > 90 % sample recovery



Ultracel® Low-binding Membranes

Fast and Efficient Concentration without Compromise


- Vertical membrane design aligns with filtrate rather than perpendicular for less clogging, less waste, and faster filtration
- Ultra-fast sample processing achieving concentration in as little as 10 minutes
- 25- to 80-fold concentration in a single step

Broad Chemical Compatibility

- Heat-sealed membrane eliminates adhesives and downstream extractables
- Large spectrum of compatibility
- Compatible with pH 1 to 9


Reliable Samples

- Spin precious samples with confidence in one robust, sleek unit that prevents leakage



Amicon® Ultra 4 mL Filters – Fast Spin Times with Excellent Recovery

Amicon® Ultra 4 mL filters were tested for percent recovery and spin time.



Consistently high recovery of diverse proteins with Amicon® Ultra filters

Concentration and percent recovery using Amicon® Ultra Filters: 4 different devices (Amicon® Ultra-0.5 mL, Amicon® Ultra-2 mL, Amicon® Ultra-4 mL, Amicon® Ultra-15 mL devices) were tested (3 kDa membrane with Cytochrome C, 10 kDa membrane with Cytochrome C, 30 kDa membrane with BSA and 100 kDa membrane with IgG) to determine percent recovery and concentration factor.

To select an Amicon® Ultra Centrifugal Filter, identify the starting volume, molecular weight of protein or nucleic acid being concentrated, final volume and concentration factor.

Then consult the product selection chart below to choose the Amicon® Ultra filter with the right nominal molecular weight cutoff (NMWCO).


| | Starting Volume | < 0.5 mL | < 2 mL | < 4 mL | < 15 mL | |
|--------------------------------|--|------------|---------|---------|---------|--|
| Molecular Weight (MW) | Proteins | | | | | |
| | | NMWCO (Da) | | | | |
| | 6 < MW < 20 k | 3,000 | 3,000 | 3,000 | 3,000 | |
| | 20 < MW < 60 k | 10,000 | 10,000 | 10,000 | 10,000 | |
| | 60 < MW < 100 k | 30,000 | 30,000 | 30,000 | 30,000 | |
| | 100 < MW < 200 k | 50,000 | 50,000 | 50,000 | 50,000 | |
| 200 k < MW | 100,000 | 100,000 | 100,000 | 100,000 | | |
| Length | Single-Stranded and Double-Stranded Nucleic Acids | | | | | |
| | | NMWCO (Da) | | | | |
| 137-1159 bp | 30,000 | 30,000 | 30,000 | 30,000 | | |
| Particle Diameter (DIA) | Nanoparticles | | | | | |
| | | NMWCO (Da) | | | | |
| | 1.5 < dia < 3 nm | 3,000 | 3,000 | 3,000 | 3,000 | |
| | 3 < dia < 5 nm | 10,000 | 10,000 | 10,000 | 10,000 | |
| | 5 < dia < 7 nm | 30,000 | 30,000 | 30,000 | 30,000 | |
| | 7 < dia < 10 nm | 50,000 | 50,000 | 50,000 | 50,000 | |
| 10 nm < dia | 100,000 | 100,000 | 100,000 | 100,000 | | |

NMWCO: Nominal Molecular Weight Cut Off


10,000 NMWCO Amicon® Ultra-4 and -15 filters are both marked and registered for *in vitro* diagnostic use.

Once you've chosen the right Amicon® Ultra filter for your needs, choose your rotor, G force and spinning time for concentrating your molecule. Designed as standard 1.5 mL, 15 mL conical or 50 mL conical tubes, Amicon® Ultra filters fit all standard rotor types.


**Amicon®
Ultra-0.5
filter**




**Amicon®
Ultra-2
filter**



**Amicon®
Ultra-4
filter**



**Amicon®
Ultra-15
filter**



| | Starting Volume | < 0.5 mL | < 2 mL | < 4 mL | < 15 mL |
|---|--------------------------|----------------------------------|---|--|----------------------------------|
| Choose a rotor and G force | Final Volume | 15–20 µL | 15–70 µL | 50 µL | 200 µL |
| | Design of the Device | Standard 1.5 mL | Standard 15 mL | Standard 15 mL | Standard 50 mL |
| | Fixed-Angle (35 °) Rotor | 14,000 g 1,000 g reverse spin | 7,500 g 1,000 g reverse spin | 5,000 g for 100,000 7,500 g for all other MWCO | 5,000 g |
| | Swinging Bucket Rotor | N/A | 4,000 g 1,000 g reverse spin | 4,000 g | 4,000 g |
| Concentration Factor | Final Volume | 15–20 µL with reverse spin | 15–70 µL with reverse spin | 50 µL | 200 µL |
| | Concentration Factor | X25–X30 | X14–X67 | X80 | X75 |
| For Proteins and Nanoparticles | | | | | |
| Adjust spinning time | NMWCO (Da) | | | | |
| | 3,000 | 30 min. | 60 min. | 40 min. | 40 min. |
| | 10,000 | 15 min. | 40 min. | 15 min. | 20 min. |
| | 30,000 | 10 min. | 20 min. | 10 min. | 20 min. |
| | 50,000 | 10 min. | 15 min. | 10 min. | 15 min. |
| | 100,000 | 10 min. | 30 min. | 10 min. | 15 min. |
| Single-Stranded and Double-Stranded Nucleic Acids | | | | | |
| | 30,000 | 10 min. | 15 min., fixed angle 40 min., swinging rotor | 10 min., 5,000 g, fixed angle | 10 min., 5,000 g, fixed angle |

Amicon® Ultra Centrifugal Filters

| | Product | Amicon® Ultra-0.5 filter | Amicon® Ultra-2 filter | Amicon® Ultra-4 filter | Amicon® Ultra-15 filter |
|------------|---|--------------------------|------------------------|------------------------|-------------------------|
| | Maximum initial sample volume (mL) | 0.5 | 2 | 4 | 15 |
| | Final concentrate (retentate) volume (µL) | 15-20 | 15-70 | 30-70 | 150-300 |
| NMWCO (Da) | Qty/Pk | | | | |
| 3,000 | 8 | UFC500308 | | UFC800308 | UFC900308 |
| | 24 | UFC500324 | UFC200324 | UFC800324 | UFC900324 |
| | 96 | UFC500396 | | UFC800396 | UFC900396 |
| | 500 | UFC5003BK | | | |
| 10,000 | 8 | UFC501008 | | UFC801008 | UFC901008 |
| | 24 | UFC501024 | UFC201024 | UFC801024 | UFC901024 |
| | 96 | UFC501096 | | UFC801096 | UFC901096 |
| | 500 | UFC5010BK | | | |
| 30,000 | 8 | UFC503008 | | UFC803008 | UFC903008 |
| | 24 | UFC503024 | UFC203024 | UFC803024 | UFC903024 |
| | 96 | UFC503096 | | UFC803096 | UFC903096 |
| | 500 | UFC5030BK | | | |
| 50,000 | 8 | UFC505008 | | UFC805008 | UFC905008 |
| | 24 | UFC505024 | UFC205024 | UFC805024 | UFC905024 |
| | 96 | UFC505096 | | UFC805096 | UFC905096 |
| | 500 | UFC5050BK | | | |
| 100,000 | 8 | UFC510008 | | UFC810008 | UFC910008 |
| | 24 | UFC510024 | UFC210024 | UFC810024 | UFC910024 |
| | 96 | UFC510096 | | UFC810096 | UFC910096 |
| | 500 | UFC5100BK | | | |

Amicon® Ultra-4 and -15 Centrifugal Filters, registered for IVD use

| Description | NMWCO | Qty/Pk | Catalog No. |
|-------------------------------------|--------|--------|-------------|
| Amicon® Ultra-4 Centrifugal Filter | 10 KDa | 8 | UFC801008D |
| | | 24 | UFC801024D |
| | | 96 | UFC801096D |
| Amicon® Ultra-15 Centrifugal Filter | 10 KDa | 8 | UFC901008D |
| | | 24 | UFC901024D |
| | | 96 | UFC901096D |



Specialized Concentration Devices

Microcon® DNA Fast Flow Filter

Concentration of gDNA and Protein

Optimized for the concentration and recovery of genomic DNA with SDS buffer. The low nonspecific binding characteristics of the membrane and the other device components, coupled with its medical-grade o-ring seal, allow the device to accommodate several wash steps with minimal sample loss.



Microcon® DNA Fast Flow Filter Advantages:

- High recovery for small volumes with reverse spin (concentration factor < 20X)
- Low-binding Ultracel® membrane
- Fast processing

Microcon® Centrifugal Filters

Simply and efficiently concentrate and desalt solutions of any macromolecule with the low-binding Ultracel® membrane, using any centrifuge that can accept 1.5 mL tubes.

Microcon® Filter Advantages:

- Dual-cycle EtO treatment on the Microcon® PCR Grade Filter has been shown to render contaminating DNA unamplifiable
- Typical recoveries of > 95%, even for dilute solutions
- Reverse spin to maximize recovery, even in the smallest samples
- Convenient storage of filtrate or concentrated sample in standard microfuge tube
- Concentration factors up to 100X

Application Guidelines

| Application | Microcon® Device | | |
|---|------------------|-----|---------------|
| | 10K | 30K | DNA Fast Flow |
| Peptide and growth factor concentration | • | | |
| Protein concentration and desalting of columns eluates | • | • | |
| Protein concentration before electrophoresis or other assays | • | • | |
| Protein removal prior to HPLC | • | • | |
| Purification of macromolecular components found in tissue culture extracts and cell lysates | • | • | |
| Concentration of biological samples (antigens, antibodies, enzymes) | | • | |
| Concentration of gDNA with or without SDS buffer | | • | • |
| Concentration and desalting of nucleic acids (single- or double-stranded) | • | • | • |
| Removal of labeled nucleotides | • | • | • |
| Removal of labeled amino acids | • | • | • |
| Removal of primers from amplified DNA | | • | • |
| Removal of linkers prior to cloning | | • | • |

Ordering Information

| Description | Volume, mL | Min. final concentrate volume, µL | Qty/Pk | Catalog No. |
|--|------------|-----------------------------------|--------|-------------|
| Microcon® filter, Ultracel®-10 membrane, 10 kDa | 0.5 | 5-50 | 100 | MRCPRT010 |
| Microcon® filter, Ultracel®-30 membrane, 30 kDa | 0.5 | 5-50 | 100 | MRCF0R030 |
| Microcon® DNA Fast Flow Centrifugal Filter with Ultracel® membrane | 0.5 | 5-50 | 100 | MRCF0R100 |
| Microcon® DNA Fast Flow PCR Grade filter with Ultracel® membrane, dual cycle EtO treated | 0.5 | 5-50 | 20 | MRCF0R100ET |

Ultrafree® spin filters for clarification, filtration, and sterilization

Ultrafree®-MC and Ultrafree®-CL centrifugal filters remove particles and precipitates from aqueous and some solvent-based samples. These fast filtration units provide highly reproducible performance for sample recovery. Ultrafree® centrifugal filters are ideal for use in protein and nucleic acid solutions.

Ultrafree®-MC filter advantages:

- High recovery Durapore® (PVDF) and hydrophilic PTFE membranes
- Five different pore sizes from 0.1 to 5.0 µm
- Pre-sterilized units also available
- Fast filtration and highly reproducible performance
- Use in fixed-angle rotors for 1.5 mL tubes

Ultrafree®-CL filter advantages:

- High recovery Durapore® (PVDF) and hydrophilic PTFE membranes
- Five different pore sizes from 0.1 to 5.0 µm
- Pre-sterilized units also available
- Fast filtration and highly reproducible performance
- Use in fixed-angle rotors for 15 mL tubes



Sterile Ultrafree®-MC and CL centrifugal filter units with microporous membrane

- Easy, pre-sterilized, centrifugal sample clarification units for either 0.5 mL (MC) or 2 mL (CL) maximum volumes
- High recovery Durapore® (PVDF) membrane
- Fast filtration and highly reproducible performance
- Use in fixed-angle rotors for 1.5 mL tubes (MC) or 15 mL tubes (CL)

Ordering Information

| | Pore Size (µm) | Color | Sterility | Qty/Pk | Catalog No. |
|--|----------------|-------------|-------------|-------------|-------------|
| Filter Units with Microporous Durapore® PVDF Membrane | | | | | |
| Ultrafree®-MC Filter | 0.1 | Orange | Non-sterile | 25 | UFC30VV25 |
| | | | | 100 | UFC30VV00 |
| | | | | 250 | UFC30GVNB |
| | 0.22 | Yellow | Non-sterile | 25 | UFC30GV25 |
| | | | | 100 | UFC30GV00 |
| | | | | 250 | UFC30GVNB |
| | | | | 50 (5 x 10) | UFC30GV0S |
| | | | | 25 | UFC30HV25 |
| | | | | 100 | UFC30HV00 |
| | 0.45 | Red | Non-sterile | 25 | UFC30HV25 |
| | | | | 100 | UFC30HV00 |
| | | | | 250 | UFC30HVN0 |
| 0.65 | Purple | Non-sterile | 25 | UFC30DV25 | |
| | | | 100 | UFC30DV00 | |
| | | | 50 (5 x 10) | UFC30DV0S | |
| 6 | Dark Green | Non-sterile | 100 | UFC30SV00 | |
| Ultrafree®-CL Filter | 0.1 | Orange | Non-sterile | 25 | UFC40VV25 |
| | | | | 100 | UFC40VV00 |
| | | | | 250 | UFC40GVNB |
| | 0.22 | Yellow | Non-sterile | 25 | UFC40GV25 |
| | | | | 100 | UFC40GV00 |
| | | | | 250 | UFC40GVNB |
| | | | | 50 (5 x 10) | UFC40GV0S |
| | | | | 25 | UFC40HV25 |
| | | | | 100 | UFC40HV00 |
| | 0.45 | Red | Non-sterile | 25 | UFC40HV25 |
| | | | | 100 | UFC40HV00 |
| | | | | 250 | UFC40HVN0 |
| 0.65 | Purple | UFC4DV25 | 25 | UFC40DV25 | |
| 5 | Dark Green | UFC40SV25 | 25 | UFC40SV25 | |
| Filter Units with Microporous Hydrophilic PTFE Membrane | | | | | |
| Ultrafree®-MC Filter | 0.22 | Yellow | Non-sterile | 25 | UFC30LG25 |
| | 0.45 | Red | Non-sterile | 25 | UFC30LH25 |
| Ultrafree®-CL Filter | 0.22 | Yellow | Non-sterile | 25 | UFC40LG25 |
| | 0.45 | Red | Non-sterile | 25 | UFC40LH25 |

Concentrate high solute samples with Centriprep® filters

Centriprep® centrifugal filters are disposable ultrafiltration devices used for purifying, concentrating, and desalting biological samples (2–15 mL volume range) and for filtration applications. Offering a high flow rate, these filters come complete and are easy to use with a twist-lock cap, a filtrate collector containing a low adsorptive Ultracel® regenerated cellulose membrane, plus an air-seal cap for sample isolation.

Centriprep® filter advantages and applications:

- Unique inverse flow mode of operation with large deadstop
- Concentrate and purify particle-laden solutions of high concentrations with Ultracel® membrane
- Fast sample processing
- Fits standard swinging-bucket rotor for 50 mL tubes
- Concentrate and purify particle-laden solutions or high concentrations
- Separate low MW solutes from fermentation broths, cell culture media, cell lysates



Ordering Information

| Description | NMWCO (kDa) | Volume (mL) | Min. final concentrate volume (µL) | Qty/Pk | Catalog No. |
|--|-------------|-------------|------------------------------------|--------|-------------|
| Centriprep® filter, Ultracel®-3 membrane | 3 | 15 | 700 | 24 | 4302 |
| | | | | 96 | 4303 |
| Centriprep® filter, Ultracel®-10 membrane* | 10 | 15 | 700 | 24 | 4304 |
| | | | | 96 | 4305 |
| Centriprep® filter, Ultracel®-30 membrane* | 30 | 15 | 700 | 24 | 4306 |
| | | | | 96 | 4307 |
| Centriprep® filter, Ultracel®-50 membrane | 50 | 15 | 700 | 24 | 4310 |
| | | | | 96 | 4311 |

* Centriprep® centrifugal filter devices with Ultracel® 10K and 30K membranes are registered for *in vitro* diagnostic use.

Clinical Ultrafiltration

Separate free from protein-bound solute with Centrifree® filters

The Centrifree® filter was designed with the clinical laboratory in mind. These devices rapidly and efficiently separate free from protein-bound microsolutes in small volumes (0.15–1.0 mL) of serum, plasma, and other biological samples using ultrafiltration. Accurate partitioning occurs in minutes without dilution, change in physiologic pH, ion composition, or unbound microsolite concentration. These devices contain low-adsorptive hydrophilic membranes and O-rings without plasticizers to ensure excellent recovery.

Centrifree® filter advantages and applications:

- Separation of free from bound microsolite in serum, plasma, and other biological samples
- Determine free therapeutic drugs, testosterone, thyroxin, etc.
- Binding studies
- New drug investigations
- Deproteinization



Ordering Information

| Description | Volume, mL | Min. final concentrate volume, µL | MWCO (kDa) | Qty/Pk | Catalog No. |
|---|------------|-----------------------------------|------------|--------|-------------|
| Centrifree® Ultrafiltration device with Ultracel® PL membrane | 1 | 50 | 30 | 50 | 4104 |

Centrifree® filters are registered for *in vitro* diagnostic use.

Minicon® filters to Concentrate Multiple Clinical Samples

Minicon® concentrators are non-sterile, disposable, multiwell ultrafiltration devices designed for concentrating macromolecules in clinical specimens such as urine, cerebrospinal fluid (CSF) or other biological solutions. The concentrators, which require no additional equipment and can be operated unattended, are used by researchers and clinical laboratories worldwide as a preparatory step to increase the sensitivity of subsequent tests.

Minicon® concentrator advantages and applications:

- Concentrate urine and cerebrospinal fluid to intensify proteins that indicate abnormal or pathological states prior to analysis by electrophoresis or immunoelectrophoresis (e.g., Bence Jones proteins in urine)
- Static concentrator, requiring no accessories
- Absorbent pulls solvent and salts through ultrafilter, concentrating sample



Ordering Information

| Description | Volume, mL | Min. final concentrate volume, µL | MWCO (kDa) | Qty/Pk | Catalog No. |
|---|------------|-----------------------------------|------------|--------|-------------|
| Minicon® B15 concentrator, 8 cells/unit | 5 | 50 | 15 | 40 | 9031 |
| Minicon® CS15 concentrator, 10 cells/unit | 2.5 | 30 | 15 | 50 | 9051 |

Minicon® filters are registered for *in vitro* diagnostic use.

Large Volume Concentration

Concentration of proteins and viruses ●

The Centricon® Plus-70 centrifugal filter is designed for rapid processing of aqueous biological solutions in volumes ranging from 15 to 70 mL. Centricon® filters concentrate most 70 mL solutions down to 350 µL in as little as 25 minutes. Samples are typically concentrated in the 50X to 200X range, depending on the sample type and starting sample volume. These units are a convenient alternative to dialysis, lyophilization, or precipitation techniques.


Centricon® Plus-70 filter advantages and applications

- > 90% typical recovery
- Low hold-up volume
- Polypropylene housing minimizes binding
- True dead stop prevents spinning to dryness
- Concentrating and desalting chromatography column eluates
- Concentrating monoclonal antibodies
- Concentrating proteins or viruses from culture supernatants
- Clarifying tissue homogenates and cell lysates



Performance

Spin time with respect to filtrate volume (Ultracel® PL-30 membrane at 3500 xg)



Ordering Information

| Description | Volume, mL | Min. final concentrate volume, µL | NMWC0 | Qty/Pk | Catalog No. |
|--------------------------------|------------|-----------------------------------|-------|--------|-------------|
| Centricon® Plus-70 3K filter | 70 | 350 | 3 | 8 | UFC700308 |
| Centricon® Plus-70 10K filter | 70 | 350 | 10 | 8 | UFC701008 |
| Centricon® Plus-70 30K filter | 70 | 350 | 30 | 8 | UFC703008 |
| Centricon® Plus-70 100K filter | 70 | 330 | 100 | 8 | UFC710008 |

Amicon® Stirred Cells ●

50 mL to 400 mL concentration

Amicon® stirred cells provide high flow rates with solutions up to 10% macrosolute concentration and are capable of rapid concentration, or salt removal followed by concentration in the same unit. For protein concentration, gas pressure is applied directly to ultrafiltration cell. Solutes above the membrane's nominal molecular weight cut-off (NMWCO) are retained in cell, while water and solutes below the cut-off pass into the filtrate and out of cell.

Advantages

- Gentle magnetic stirring minimizes concentration polarization and shear denaturation.
- All stirred cells can be autoclaved.
- Three different sizes to handle volumes from 50 mL to 400 mL
- High flow rates with solutions up to 10% macrosolute concentration

Applications

- Concentrate, diafilter, and exchange buffers for macromolecule solutions including proteins, enzymes, antibodies and viruses.

Available in three sizes

| Max. Working Volume | Catalog No. |
|---------------------|-------------|
| 50 mL | UFSC05001 |
| 200 mL | UFSC20001 |
| 400 mL | UFSC40001 |



Introducing the new Amicon® Stirred Cell.

Order the new 50 mL, 200 mL, or 400 mL stirred cells and you will experience the same performance to which you're accustomed: gentle, high recovery of macrosolutes, thorough buffer exchange, membrane flexibility and ability to monitor filtration progress. In addition, you will enjoy many workflow-enhancing features.

What's new about the updated stirred cells:

- Ergonomic benefits: you will love how easy it is to open, close and assemble the new Amicon® stirred cell!
- Quick connectors to tubing for easy, secure setup.
- Integrated safety features: with screw threads and a pressure relief valve, there's no need for external housing. This means easier assembly and disassembly, and very clear confirmation that the device is properly assembled.
- Overall superior integrity (no leaking).
- Broader selection of membrane discs.
- Fully revised user guide with clearer instructions for operation and how to connect to your gas source.
- Better spare part and accessory support.
- More secure stir bar eliminates risk of damage to your membrane.

Membrane Discs for Use in Stirred Cells

Ultracel® cut disc membranes

To concentrate or desalt dilute solutions, use Ultracel® regenerated cellulose membranes. The hydrophilic, tight microstructure of Ultracel® membranes assures the highest possible retention with the lowest possible adsorption of protein, DNA or other macromolecules.

- Membranes available in 1, 3, 5, 10, 30 and 100 kDa nominal molecular weight limit (NMWL).
- Filter diameters available in 25, 44.5, 47, 63.5, 76, 90 and 150 mm.

For ordering information, visit SigmaAldrich.com/UltracelUFcutdiscs

Biomax® cut disc membranes

To concentrate or desalt higher volumes of more concentrated samples (recommended for protein concentrations greater than 1.0 µg/mL), use Biomax® polyethersulfone (PES) membranes. These membranes are recommended for samples such as serum, plasma, or conditioned tissue culture media.

- Membranes available in 5, 10, 30, 50, 100, 300, and 500 kDa nominal molecular weight limit (NMWL).
- Filter diameters available in 25, 44.5, 47, 63.5, 76, 90 and 150 mm.


For ordering information, please visit SigmaAldrich.com/BiomaxUFcutdiscs

Durapore® cut disc membranes


For large-volume microfiltration, choose Durapore® PVDF membrane discs for your stirred cell.

- Membranes available in 0.1, 0.2 and 0.45 µm pore sizes
- Filter diameters available in 63.5 and 70 mm.


Contact customer service for Durapore® discs ordering information.



Ultracel® regenerated cellulose ultrafiltration membrane.



Biomax® polyethersulfone ultrafiltration membrane.



Durapore® polyvinylidene fluoride ultrafiltration membrane.

Stirred Cell Accessories Expand Your Capabilities

Amicon® Stirred Cell Selector Valve (Catalog No. 6003)


Valve with sliding control for instant switching from concentration to diafiltration, or switching gas and liquid lines simultaneously. Simplifies operation and avoids the need for multiple T-fittings and valves.

Amicon® Stirred Cell Manifold (Catalog No. 6015)

For instant direction of gas pressure or liquid flow in multi-cell or multi-reservoir systems. Can pressurize up to 3 cells or reservoirs from one gas source or feed several cells from one reservoir.

Amicon® Stirred Cell Reservoir (Catalog No. 6028)

This 800 mL auxiliary reservoir increases the volume capacity of stirred cells. When pressurized from an external gas source, it automatically replenishes liquid in the cell's built-in reservoir during filtration. The reservoir may also be used to store dialysate during diafiltration or dialysis.



Amicon® Stirred Cell setup for large volume concentration and continuous diafiltration using selector valve and reservoir accessories.

To place an order or receive technical assistance

Order/Customer Service: [SigmaAldrich.com/order](https://www.sigmaaldrich.com/order)

Technical Service: [SigmaAldrich.com/techservice](https://www.sigmaaldrich.com/techservice)

Safety-related Information: [SigmaAldrich.com/safetycenter](https://www.sigmaaldrich.com/safetycenter)

SigmaAldrich.com

