


# A New Class of NHC-Based Pd Complexes for Challenging Cross-Couplings of Aryl Chlorides

Yuanzhe Zhang<sup>1</sup>, Meenal Kaushal<sup>1</sup>, Yam N. Timsina<sup>1</sup>, Ashish Singh<sup>1</sup>, Thomas J. Colacot<sup>1</sup>, and Guolin Xu<sup>1</sup>

<sup>1</sup>Research and Development, Life Science Chemistry, Merck

A class of N-heterocyclic carbene (NHC)-based palladates,  $(\text{NHC-H})_2\text{Pd}_2\text{Cl}_6$  (NHC = IPr, SIPr, IPent, IPr\*, IPr\*OMe and IPr#), was synthesized in excellent yields and characterized. Comparison of their catalytic activities in Suzuki-Miyaura coupling with other NHC-Pd-based precatalysts revealed comparable or


superior activities, besides avoiding the use of “throw-away” stabilizing ligands. These precatalysts were demonstrated for Suzuki-Miyaura and Buchwald-Hartwig couplings with good substrate scope and isolated yields. In addition, transamidation coupling was also demonstrated.



**Figure 1.** Structure, synthesis, and catalytic applications of  $(\text{NHC-H})_2\text{Pd}_2\text{Cl}_6$  precatalysts. The schematic shows a plausible activation pathway of  $(\text{NHC-H})_2\text{Pd}_2\text{Cl}_6$  precatalysts prior to the cross-coupling catalytic cycle. Applications include Suzuki-Miyaura, Buchwald-Hartwig, and transamidation reactions. Highlights note scalable synthesis and broad utility.

## Past Solutions, Present Challenges in Pd–NHC Catalysis

N-heterocyclic carbenes (NHCs) have emerged as a distinctive and powerful class of ligands in cross-coupling chemistry, particularly for the development of palladium-based precatalysts.<sup>1–15</sup> The majority of the studies in this area were focused on palladium-catalyzed processes with significant contributions from the groups of Nolan,<sup>10,16–18</sup> Organ,<sup>19,20</sup> Glorius,<sup>15</sup> Szostak,<sup>21–24</sup> Hazari,<sup>16,25–28</sup> and others<sup>29–33</sup> (Figure 2). To stabilize the highly reactive 12-electron NHC–Pd(0) species formed during catalysis, chemists have adopted the use of “throw-away” stabilizing ligands such as allyl, crotyl, cinnamyl, 3-chloropyridyl, and t-butyl indenyl. However, incorporating these ligands onto palladium remains a nontrivial challenge.<sup>1</sup>



**Figure 2.** Examples of reported NHC–Pd precatalysts with “throw-away” stabilizing ligands (red) as reported by Nolan (2002, 2006), Organ (2006), Hazari (2015), and Szostak (2021). Image shows four Pd–NHC precatalysts with different, red-highlighted stabilizing ligands, emphasizing their temporary role in catalysis.

Among modern NHC-based catalytic systems involving monoligated Pd–NHC complexes with spectator/throw-away ligands, a notable breakthrough came in 2002 when Nolan introduced a series of (NHC)Pd(allyl)Cl precatalysts stabilized by allyl groups.<sup>34</sup> Building on this, the group later developed (NHC)Pd(R-allyl)Cl complexes in 2006, featuring Me or Ph substitutions on the allyl ligand, which allowed room-temperature activation compared to the 70 °C required for the original system.<sup>35</sup> While these precatalysts were effective in cross-coupling reactions, they were prone to forming inactive bridging Pd(I)–allyl dimers during activation, negatively impacting catalytic efficiency. However, bulkier substituents on the allyl group or ligand backbone helped mitigate dimer formation and improve activation.<sup>26,36–38</sup>

## Toward Scalable and Ligand-Free NHC–Pd Precatalysts


While Pd–PEPSI precatalysts are accessible in good yields, their synthesis typically requires relatively harsh conditions (80 °C in 3-chloropyridine for 16 hours) and time-intensive workups, limiting their scalability for bulk applications.

In 2015, the Hazari group introduced a new class of NHC–Pd precatalysts stabilized by η<sup>3</sup>-1-tBu-indenyl ligands.<sup>25,28</sup> Unlike their R-allyl counterparts, these complexes were designed to suppress the formation of inactive Pd(I) dimers formed via comproportionation and demonstrated strong performance in Suzuki–Miyaura

and Buchwald–Hartwig couplings. However, large-scale synthesis remains challenging, often involving multi-step procedures and the difficult incorporation of the t-butyl group into the indenyl framework. The latter step raises particular safety concerns due to the use of tBuLi.<sup>25,40</sup> More recently, Szostak and coworkers reported a series of aniline-stabilized NHC–Pd precatalysts, (NHC)Pd(AN)Cl<sub>2</sub> (AN = aniline or m-CF<sub>3</sub> aniline), showcasing their broad utility in **cross-coupling reactions**.<sup>21,23</sup>

(The cross-coupling guide seen here is geo-targeted) Although many of these NHC–Pd-based precatalysts containing stabilizing ligands are widely utilized in both academic research and industrial applications, their limited Pd atom economy and (relatively) complex manufacturing processes restrict broader adoption due to high cost and scalability challenges. Developing a one-pot, high-yield synthesis of NHC–Pd precatalysts could significantly advance the field.

Recently, stabilizing-ligand-free NHC–Pd precatalysts have garnered attention, with contributions from Nolan, Szostak, and Gooßen (Figure 3A).<sup>24,41–45</sup> Complexes such as [Pd(NHC)(μ-Cl)Cl]<sub>2</sub> have shown promise in Suzuki–Miyaura and Buchwald–Hartwig couplings but often rely on less practical reagents like Ag<sub>2</sub>O or corrosive HCl, limiting broader applicability.<sup>24,41,43,44</sup>




**Figure 3.** Examples of stabilizing-ligand-free NHC–Pd precatalysts (A), monodentate phosphine palladates (B), and NHC–Pd precatalysts studied (C). Includes control studies on (NHC–H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> activation with key reactions (70–100% yields) and a proposed catalytic cycle showing oxidative addition, reduction, and transmetalation. Inset: 3D model of Pd–butylamine complex.

Herein, we present a new class of stabilizing-ligand-free NHC–Pd precatalysts, (NHC–H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> (Figure 3C), synthesized under mild, scalable conditions and demonstrating excellent catalytic performance. While similar complexes were reported by Trzeciak in 2015 via a two-step synthesis (60% overall yield),<sup>46,47</sup> and by Nolan in a mechanistic study,<sup>45</sup> neither group explored their application in cross-coupling. Related phosphine-based systems, such as [P–H]<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> (Figure 3B), have also been used in cross-coupling,<sup>48</sup> though they rely on bulky monodentate ligands like BrettPhos or CataCXium.

## RESULTS AND DISCUSSION: Efficient Synthesis and Characterization of (NHC-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> Complexes

Unlike the synthesis of (IPr-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> reported by the Nolan group, which required elevated temperature (60 °C) in a flammable organic solvent (acetone),<sup>45</sup> The (NHC-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> complexes reported herein were all synthesized under much milder conditions (reacting Na<sub>2</sub>PdCl<sub>4</sub> with NHC salts in methanol/water at room temperature). All products were isolated in excellent yields after a simple filtration (>80% isolated, Scheme 1). Four of these NHC palladates, namely, (IPr-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> (1a), (SIPr-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> (1b), (IPent-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> (1c), and (IPr#-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> (1f), were scaled up to >20 grams without any yield loss, demonstrating the scalability of the process for large-scale applications and consequently cost savings and sustainability.




**Scheme 1.** Synthesis of (NHC-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> (1a–f). (NHC-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> complexes were prepared by reacting NHC salts with Na<sub>2</sub>PdCl<sub>4</sub> in water/methanol at room temperature for 1 hour. Variations in NHC ligands (1a–f) yielded 80–94%, producing Pd dimers with square planar geometry coordinated by two NHC ligands and four chlorides

All the products were fully characterized with <sup>1</sup>H NMR, <sup>13</sup>C NMR and elemental analysis. (IPr-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> (1a), (SIPr-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> (1b) and (IPr#-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> (1f) were confirmed with single-crystal X-ray crystallography.

### Activation Pathways and Catalytic Performance of NHC-Pd Complexes

In principle, these complexes need to be activated cooperatively on both [NHC-H]<sup>+</sup> and [Pd<sub>2</sub>Cl<sub>6</sub>]<sup>2-</sup> motifs. We carried out two controlled reactions: i) reacting (IPr\*-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> (1d) with base (NaOtBu) first, followed by the addition of amine (nBuNH<sub>2</sub>) (reaction 1 in **Scheme 2A**); and ii) reacting (IPr\*-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> (1d) with amine first, followed by base (reaction 2 in **Scheme 2A**). Both reactions gave precatalyst 2 in good yields (reaction 1, 70%; reaction 2, 90%). With these observations, we were confident that, with the presence of an amine, our (NHC-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> precatalysts can be activated to form 2 as the active precatalyst in cross-coupling reactions. In addition, precatalyst 2 was utilized for a Buchwald-Hartwig reaction (reaction 3 in **Scheme 2A**) with quantitative conversion. Combining all these findings, we believe that these (NHC-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> precatalysts can be utilized as the effective catalysts in cross-coupling reactions with the proposed mechanism in **Scheme 2B**.




**Scheme 2.** Control studies on (NHC-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> activation mechanism (A) showing three key reactions with isolated yields of 70–100%, and (B) the proposed catalytic cycle illustrating oxidative addition, reduction, and transmetalation steps involving NHC salts and Pd intermediates. The inset features a 3D model of the activated Pd complex coordinated with butylamine and chloride.

Given the structural and mechanistic similarities between our (NHC-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> precatalysts and established systems such as the Pd-PEPPSI series, (NHC)Pd(R-allyl)Cl, and (NHC)Pd(tBu-Ind)Cl, we anticipated comparable, if not superior, performance in cross-coupling reactions. We were particularly interested in whether these commonly used “throw-away” stabilizing ligands (e.g., 3-chloropyridine, tBu-indenyl) could be entirely omitted without compromising catalytic activity.

To explore the necessity of “throw-away” stabilizing ligands, we compared our (IPent-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> (1c) with two commercial precatalysts, Pd-PEPPSI-IPent and (IPent)Pd(tBu-Ind)Cl, in a challenging Suzuki-Miyaura reaction between chloromesitylene and 2,6-dimethylphenylboronic acid (**Figure 4**). While all three achieved >90% conversion, reaction rates varied notably. Our catalyst (1c) completed the reaction in ~10 h, faster than Pd-PEPPSI-IPent (~20 h), suggesting the stabilizing 3-chloropyridine may impede turnover. The tBu-Ind-stabilized precatalyst showed the fastest rate (~6 h). While differences were moderate, these results support the idea that eliminating stabilizing ligands can maintain activity while improving atom economy and simplifying downstream purification.

## Broad Applications in Cross-Coupling Reactions


We then expanded our application studies with respect to the substrate scopes and types of cross-coupling reactions (i.e., Suzuki-Miyaura coupling, Buchwald-Hartwig coupling, and trans-amidation coupling). While cross-coupling reactions of aryl bromide catalyzed by NHC-Pd-based precatalysts were well documented in literature, we were particularly interested in the cross-coupling chemistry of relatively challenging aryl chlorides. In the Suzuki-Miyaura coupling reactions (Figure 4), using the previously demonstrated conditions (1–2 mol% Pd, KOH as base), sterically hindered aryl-aryl ( $sp^2-sp^2$ ) couplings performed



**Figure 4.2.** Substrate scope, d. aReaction conditions: (Het)Ar-Cl (0.5 mmol, 1.0 equiv.), boronic acid/ester (1.5–2 equiv.), KOH (2.0 equiv.), [NHC-H]<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> (1.0 mol% Pd), solvent; dioxane, 60 °C, GC conversion, isolated yield; d for  $sp^2-sp^3$ : (IPr<sup>\*</sup>-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> 1f (1.0 mol% Pd), solvent; dioxane, 90 °C; control; eno catalyst. Suzuki-Miyaura coupling scheme and graph comparing conversion rates of Pd catalysts, with (IPent-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> reaching 90% at 10 hours

well, with isolated yields exceeding 80%. For example, using (IPent-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> (**1c**), product **4a** was obtained in 84% yield. Similarly, (IPr<sup>\*</sup>-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> (**1d**) delivered **4a** in 80% yield and **4b** in 84%.


In contrast, for less hindered substrates such as product **4c**, (IPent-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> (**1a**) gave excellent results (81% yield), although this same precatalyst gave negligible conversion (<10%) for the more hindered **4a** and **4b**. Notably, nitrogen-containing substrates like pyridine were also well tolerated, as demonstrated by the 91% isolated yield of **4d**.



**Figure 4.1.** Suzuki-Miyaura coupling using (NHC-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> precatalysts, benchmarking (IPent-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> against Pd-PEPPSI-IPent and (IPent)Pd(tBu-Ind)Cl. Substrates with key, red-highlighted groups and yields (80–94%) show catalyst effectiveness in Suzuki-Miyaura coupling on a clear white background.

Although  $sp^2-sp^3$  cross-coupling reactions using NHC-Pd catalysts have been demonstrated with Pd-PEPPSI precatalysts in Negishi and Feringa-Murahashi systems, Suzuki-Miyaura-type  $sp^2-sp^3$  couplings remain less

explored with NHC-Pd systems.<sup>49, 51–60</sup> Nevertheless, we successfully applied our (NHC-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> catalysts in these challenging reactions, achieving strong performance within a limited substrate scope.



**Figure 5.** Buchwald-Hartwig coupling study using (NHC-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> precatalysts, showing broad substrate scope, high yields, and selective halogen reactivity. <sup>f</sup>Reaction conditions: Ar-X (1.0 mmol), amine (1.5 equiv.), NaOtBu (1.5 mmol, 1.5 equiv.), [IPr-H]<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> (**1d** 0.5 mol%), 2-MeTHF (1 mL), 55 °C; run at 105 °C; isolated as HCl salt; K<sub>3</sub>PO<sub>4</sub> (3 equiv.), dioxane, 100 °C. Table summarizing Buchwald-Hartwig coupling with (NHC-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> showing substrates, nitrogen highlights, and yields from 65% to 97%.

Using (IPr#-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> (1–2 mol% Pd loading), coupling of 4-chloroquinoline with methylboronic acid or cyclopropylboronic acid yielded products **5a** and **5b** in 92% and 86% isolated yields, respectively. No product was formed in the absence of a catalyst.


Additionally, stepwise coupling was observed for Ar-Cl with two chloride groups of differing reactivity, such as 2,6-dichloroquinoline. Mono-coupled products **5c** and **5d** were obtained in good yields (62% and 52%, respectively), with the second chloride remaining untouched.

NHC-Pd-catalyzed Buchwald-Hartwig couplings of aryl chlorides have been broadly demonstrated in literature

Our (NHC-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> precatalysts also enabled efficient cascade reactions and exhibited notable haloselectivity. Indole derivatives **6k** and **6l** were obtained in excellent yields via cascade Buchwald-Hartwig coupling of primary amines followed by hydroamination of a C–C triple bond.<sup>62</sup> Notably, the catalysts demonstrated excellent selectivity between halogens on polyhalogenated substrates. Selective amination occurred preferentially at the Br position over Cl in aryl halides (Ar-X), affording products **6m–6p** in good to excellent yields. This selectivity is largely due to the reactivity difference between the Br and Cl groups on the substrates. Similarly, product **6q** was obtained successfully from a selective coupling reaction on the relatively more active Cl group of 2,6-dichloroquinoline and cyclobutylamine (**6q**, 92%).

with significant contributions from the groups of Nolan,<sup>17,34,35</sup> Organ,<sup>20,50</sup> Hazari,<sup>18,25,28</sup> Szostak,<sup>21,22,44,58</sup> and many others.<sup>5,61</sup> Without too much surprise, (NHC-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> also worked successfully with broad substrate scopes and excellent isolated yields (**Figure 5**). Select coupling of primary amines (**6a–f**) and secondary amines (**6g** and **6h**) was generally observed. Both aryl amines and alkyl amines were successfully utilized in the coupling reactions catalyzed by (NHC-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> precatalysts. Worthwhile to note, primary hydrazones, i.e., benzophenone hydrazone, reacted with aryl chlorides smoothly, even in the case of the sterically hindered 2,6-dimethylchlorobenzene, with excellent isolated yields (**6i**, 93%; **6j**, 94%, respectively).

Lastly, we briefly expanded the applications of our (NHC-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> into Pd-catalyzed trans-amidation couplings.<sup>58</sup> Both ester trans-amidation and amide trans-amidation worked smoothly with moderate to excellent isolated yields. For instance, in **Figure 6**, product **7a** was isolated in good to excellent yields from both trans-amidation reactions (ester trans-amidation, 84%; amide trans-amidation, 91%). However, when the alkyl ester or amide was used as the electrophile, the reaction turned out to be much more sluggish with moderate isolated yields (**7b**, 62%; **7c**, 40%). Boc-protected amino acid ester was well tolerated and utilized for ester trans-amidation with good yield (**7c**, 40%).



**Figure 6.** Use of (NHC-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> precatalysts in ester and amide trans-amidation coupling, delivering moderate to excellent yields and tolerating Boc-protected substrates. **Reaction conditions:** Aniline (0.5 mmol, 2.0 equiv.), tertiary amide or Aryl benzoate (0.25 mmol, 1.0 equiv.), K<sub>2</sub>CO<sub>3</sub> (0.75 mmol, 1.5 equiv.), (IPr\*-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub>, (**1d**, 3 mol%), DME (1 mL), 100 °C.

In conclusion, we have developed and characterized a class of N-heterocyclic carbene (NHC)-based palladates, (NHC-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub> (NHC = IPr, SIPr, IPent, IPr\*, IPr\*OMe and IPr#) that operate without the need for stabilizing “throw-away” ligands. Mechanistic studies and direct benchmarking against leading commercial NHC–Pd precatalysts revealed comparable or superior

catalytic activity. These ligand-free systems performed exceptionally well across diverse and industrially relevant cross-coupling chemistries, including Suzuki–Miyaura, Buchwald–Hartwig, and trans-amidation couplings, offering broad substrate scope, excellent isolated yields, and promising potential for sustainable large-scale applications.

## REFERENCES

- Firsan, S. J.; Sivakumar, V.; Colacot, T. J. Emerging Trends in Cross-Coupling: Twelve-Electron-Based L1Pd(0) Catalysts, Their Mechanism of Action, and Selected Applications. *Chem. Rev.* **2022**, *122* (23), 16983–17027. <https://doi.org/10.1021/acs.chemrev.2c00204>.
- Fricke, C.; Sperger, T.; Mendel, M.; Schoenebeck, F. Catalysis with Palladium(I) Dimers. *Angew. Chemie - Int. Ed.* **2021**, *60* (7), 3355–3366. <https://doi.org/10.1002/anie.202011825>.
- Gildner, P. G.; Colacot, T. J. Reactions of the 21st Century: Two Decades of Innovative Catalyst Design for Palladium-Catalyzed Cross-Couplings. *Organometallics* **2015**, *34* (23), 5497–5508. <https://doi.org/10.1021/acs.organomet.5b00567>.
- Johansson Seechurn, C. C. C.; Kitching, M. O.; Colacot, T. J.; Snieckus, V. Palladium-Catalyzed Cross-Coupling: A Historical Contextual Perspective to the 2010 Nobel Prize. *Angew. Chemie - Int. Ed.* **2012**, *51* (21), 5062–5085. <https://doi.org/10.1002/anie.201107017>.
- Shaughnessy, K. H. Development of Palladium Precatalysts That Efficiently Generate LPd(0) Active Species. *Isr. J. Chem.* **2020**, *60*, 180. <https://doi.org/10.1002/ijch.201900067>.
- Bellotti, P.; Koy, M.; Hopkinson, M. N.; Glorius, F. Recent Advances in the Chemistry and Applications of N-Heterocyclic Carbenes. *Nat. Rev. Chem.* **2021**, *5* (10), 711–725. <https://doi.org/10.1038/s41570-021-00321-1>.
- Benhamou, L.; Chardon, E.; Lavigne, G.; Bellemin-Laponnaz, S.; César, V. Synthetic Routes to N-Heterocyclic Carbene Precursors. *Chem. Rev.* **2011**, *111* (4), 2705–2733. <https://doi.org/10.1021/cr100328e>.
- Enders, D.; Niemeier, O.; Henseler, A. Organocatalysis by N-Heterocyclic Carbenes. *Chem. Rev.* **2007**, *107* (12), 5606–5655. <https://doi.org/10.1021/cr068372>.
- Flanigan, D. M.; Romanov-Mikhailidis, F.; White, N. A.; Rovis, T. Organocatalytic Reactions Enabled by N-Heterocyclic Carbenes. *Chem. Rev.* **2015**, *115* (17), 9307–9387. <https://doi.org/10.1021/acs.chemrev.5b00060>.
- Fortman, G. C.; Nolan, S. P. N-Heterocyclic Carbene (NHC) Ligands and Palladium in Homogeneous Cross-Coupling Catalysis: A Perfect Union. *Chem. Soc. Rev.* **2011**, *40* (10), 5151–5169. <https://doi.org/10.1039/c1cs15088j>.
- Hopkinson, M. N.; Richter, C.; Schedler, M.; Glorius, F. An Overview of N-Heterocyclic Carbenes. *Nature*. Nature Publishing Group **2014**, pp 485–496. <https://doi.org/10.1038/nature13384>.
- Huynh, H. V. Electronic Properties of N-Heterocyclic Carbenes and Their Experimental Determination. *Chem. Rev.* **2018**, *118* (19), 9457–9492. <https://doi.org/10.1021/acs.chemrev.8b00067>.
- Lee, B. C.; Liu, C. F.; Lin, L. Q. H.; Yap, K. Z.; Song, N. X.; Ko, C. H. M.; Chan, P. H.; Koh, M. J. N-Heterocyclic Carbenes as Privileged Ligands for Nickel-Catalysed Alkene Functionalisation. *Chem. Soc. Rev.* **2023**, *52* (9), 2946–2991. <https://doi.org/10.1039/d2cs00972b>.
- Peris, E. Smart N-Heterocyclic Carbene Ligands in Catalysis. *Chem. Rev.* **2018**, *118* (19), 9988–10031. <https://doi.org/10.1021/acs.chemrev.6b00695>.
- Würtz, S.; Glorius, F. Surveying Sterically Demanding N-Heterocyclic Carbene Ligands with Restricted Flexibility for Palladium-Catalyzed Cross-Coupling Reactions. *Acc. Chem. Res.* **2008**, *41* (11), 1523–1533. <https://doi.org/10.1021/ar8000876>.
- Izquierdo, F.; Zinser, C.; Minenkov, Y.; Cordes, D. B.; Slawin, A. M. Z.; Cavallo, L.; Nahra, F.; Cazin, C. S. J.; Nolan, S. P. Insights into the Catalytic Activity of [Pd(NHC)(Cin)Cl] (NHC=IPr, IPrCl, IPrBr) Complexes in the Suzuki–Miyaura Reaction. *ChemCatChem* **2018**, *10* (3), 601–611. <https://doi.org/10.1002/cctc.201701279>.
- Marion, N.; Ecarnot, E. C.; Navarro, O.; Amoroso, D.; Bell, A.; Nolan, S. P. (IPr)Pd(AcAc)Cl: An Easily Synthesized, Efficient, and Versatile Precatalyst for C–N and C–C Bond Formation. *J. Org. Chem.* **2006**, *71* (10), 3816–3821. <https://doi.org/10.1021/jo060190h>.
- Zhou, T.; Li, G.; Nolan, S. P.; Szostak, M. Well-Defined, Air-Stable, and Readily Available Precatalysts for Suzuki and Buchwald–Hartwig Cross-Coupling (Transamidation) of Amides and Esters by N–C/O–C Activation. *Org. Lett.* **2019**, *21* (9), 3304–3309. <https://doi.org/10.1021/acs.orglett.9b01053>.
- Kantchev, E. A. B.; O'Brien, C. J.; Organ, M. G. *Palladium Complexes of N-Heterocyclic Carbenes as Catalysts for Cross-Coupling Reactions - A Synthetic Chemist's Perspective*; **2007**; Vol. 46. <https://doi.org/10.1002/anie.200601663>.
- (20) Froese, R. D. J.; Lombardi, C.; Pompeo, M.; Rucker, R. P.; Organ, M. G. Designing Pd–N-Heterocyclic Carbene Complexes for High Reactivity and Selectivity for Cross-Coupling Applications. *Acc. Chem. Res.* **2017**, *50* (9), 2244–2253. <https://doi.org/10.1021/acs.accounts.7b00249>.
- Xia, Q.; Shi, S.; Gao, P.; Lalancette, R.; Szostak, R.; Szostak, M. Complexes: Easily Synthesized, Highly Active Pd(II)-NHC Precatalysts for Cross-Coupling Reactions. *J. Org. Chem.* **2021**, *86* (21), 15648–15657. <https://doi.org/10.1021/acs.joc.1c02183>.
- Li, D. H.; Lan, X. B.; Song, A. X.; Rahman, M. M.; Xu, C.; Huang, F. D.; Szostak, R.; Szostak, M.; Liu, F. S. Buchwald–Hartwig Amination of Coordinating Heterocycles Enabled by Large-but-Flexible Pd–BIAN–NHC Catalysts. *Chem. - A Eur. J.* **2022**, *28* (4), e202103341. <https://doi.org/10.1002/chem.202103341>.
- Lei, P.; Wang, Y.; Zhang, C.; Hu, Y.; Feng, J.; Ma, Z.; Liu, X.; Szostak, R.; Szostak, M. Sonogashira Cross-Coupling of Aryl Ammonium Salts by Selective C–N Activation Catalyzed by Air- and Moisture-Stable, Highly Active [Pd(NHC)(3-CF<sub>3</sub>-An)Cl<sub>2</sub>] (An = Aniline) Precatalysts. *Org. Lett.* **2022**, *24*, 6310–6315. <https://doi.org/10.1021/acs.orglett.2c02534>.
- Yang, S.; Li, H.; Yu, X.; An, J.; Szostak, M. Suzuki–Miyaura Cross-Coupling of Aryl Fluorosulfonates Mediated by Air- and Moisture-Stable [Pd(NHC)(μ-Cl)Cl]<sub>2</sub>Precatalysts: Broad Platform for C–O Cross-Coupling of Stable Phenolic Electrophiles. *J. Org. Chem.* **2022**, *87* (22), 15250–15260. <https://doi.org/10.1021/acs.joc.2c01778>.
- Melvin, P. R.; Nova, A.; Balcells, D.; Dai, W.; Hazari, N.; Hruszkewycz, D. P.; Shah, H. P.; Tudge, M. T. Design of a Versatile and Improved Precatalyst Scaffold for Palladium-Catalyzed Cross-Coupling: (H<sub>3</sub>-1-TBu-Indenyl)<sub>2</sub>(μ-Cl)<sub>2</sub>Pd<sub>2</sub>. *ACS Catal.* **2015**, *5* (6), 3680–3688. <https://doi.org/10.1021/acscatal.5b00878>.
- Melvin, P. R.; Balcells, D.; Hazari, N.; Nova, A. Understanding Precatalyst Activation in Cross-Coupling Reactions: Alcohol Facilitated Reduction from Pd(II) to Pd(0) in Precatalysts of the Type (H<sub>3</sub>-Allyl)Pd(L)(Cl) and (H<sub>3</sub>-Indenyl)Pd(L)(Cl). *ACS Catal.* **2015**, *5* (9), 5596–5606. <https://doi.org/10.1021/acscatal.5b01291>.
- Hazari, N.; Melvin, P. R.; Beromi, M. M. Well-Defined Nickel and Palladium Precatalysts for Cross-Coupling. *Nat. Rev. Chem.* **2017**, *1* (3), 1–17. <https://doi.org/10.1038/s41570-017-0025>.
- Dardir, A. H.; Melvin, P. R.; Davis, R. M.; Hazari, N.; Mohadjer Beromi, M. Rapidly Activating Pd–Precatalyst for Suzuki–Miyaura and Buchwald–Hartwig Couplings of Aryl Esters. *J. Org. Chem.* **2018**, *83* (1), 469–477. <https://doi.org/10.1021/acs.joc.7b02588>.
- Titcomb, L. R.; Caddick, S.; Geoffrey, F.; Cloke, N.; Wilson, D. J.; McKerrecher, D. Unexpected Reactivity of Two-Coordinate Palladium–Carbene Complexes; Synthetic and Catalytic Implications. *Chem. Commun.* **2001**, No. 15, 1388–1389. <https://doi.org/10.1039/b104297c>.
- Lewis, A. K. d. K.; Caddick, S.; Cloke, F. G. N.; Billingham, N. C.; Hitchcock, P. B.; Leonard, J. Synthetic, Structural, and Mechanistic Studies on the Oxidative Addition of Aromatic Chlorides to a Palladium (n-Heterocyclic Carbene) Complex: Relevance to Catalytic Amination. *J. Am. Chem. Soc.* **2003**, *125* (33), 10066–10073. <https://doi.org/10.1021/ja035565k>.
- César, V.; Bellemin-Laponnaz, S.; Gade, L. H. Direct Coupling of Oxazolines and N-Heterocyclic Carbenes: A Modular Approach to a New Class of C–N Donor Ligands for Homogeneous Catalysis. *Organometallics* **2002**, *21* (24), 5204–5208. <https://doi.org/10.1021/om020608b>.
- R. Jackstell, M. G. Andreu, A. Frisch, K. Selvakumar, A. Zapf, H. Klein, A. Spannenberg, D. Rottger, O. Briel, R. Karch, M. B. A Highly Efficient Catalyst for the Telomerization of 1,3-Dienes with Alcohols: First Synthesis of a Monocarbene-palladium(0) $\pm$  Olefin Complex. *Angew. Chem. Int. Ed.* **2002**, *41* (6), 12–26.
- Schneider, S. K.; Herrmann, W. A.; Herdtweck, E. Active Catalysts for the Suzuki Coupling: Palladium Complexes of Tetrahydropyrimid-2-Ylidenes. *J. Mol. Catal. A Chem.* **2006**, *245* (1–2), 248–254. <https://doi.org/10.1016/j.molcata.2005.08.046>.
- Viciu, M. S.; Germaneau, R. F.; Navarro-Fernandez, O.; Stevens, E. D.; Nolan, S. P. Activation and Reactivity of (NHC)Pd(Allyl)

- Cl (NHC = N-Heterocyclic Carbene) Complexes in Cross-Coupling Reactions. *Organometallics* **2002**, 21 (25), 5470–5472. <https://doi.org/10.1021/om020804i>.
35. Marion, N.; Navarro, O.; Mei, J.; Stevens, E. D.; Scott, N. M.; Nolan, S. P. Modified ( NHC ) Pd ( Allyl ) Cl ( NHC ) N-Heterocyclic Carbene ) Complexes for Room-Temperature Suzuki - Miyaura and Buchwald - Hartwig Reactions. **2006**, No. 11, 4101–4111.
  36. Deangelis, A. J.; Gildner, P. G.; Chow, R.; Colacot, T. J. Generating Active L-Pd(0) via Neutral or Cationic n-Allylpalladium Complexes Featuring Biaryl/Bipyrazolylphosphines: Synthetic, Mechanistic, and Structure-Activity Studies in Challenging Cross-Coupling Reactions. *J. Org. Chem.* **2015**, 80 (13), 6794–6813. <https://doi.org/10.1021/acs.joc.5b01005>.
  37. Hill, L. L.; Crowell, J. L.; Tutwiler, S. L.; Massie, N. L.; Hines, C. C.; Griffin, S. T.; Rogers, R. D.; Shaughnessy, K. H.; Grasa, G. A.; Johansson Seechurn, C. C. C.; Li, H.; Colacot, T. J.; Chou, J.; Woltermann, C. J. Synthesis and X-Ray Structure Determination of Highly Active Pd(II), Pd(I), and Pd(0) Complexes of Di(Tert -Butyl) Neopentylphosphine (DTBnP) in the Arylation of Amines and Ketones. *J. Org. Chem.* **2010**, 75 (19), 6477–6488. <https://doi.org/10.1021/jo101187q>.
  38. Hruszkewycz, D. P.; Balcells, D.; Guard, L. M.; Hazari, N.; Tilset, M. Insight into the Efficiency of Cinnamyl-Supported Precatalysts for the Suzuki-Miyaura Reaction: Observation of Pd(I) Dimers with Bridging Allyl Ligands during Catalysis. *J. Am. Chem. Soc.* **2014**, 136 (20), 7300–7316. <https://doi.org/10.1021/ja412565c>.
  39. O'Brien, C. J.; Kantchev, E. A. B.; Valente, C.; Hadei, N.; Chass, G. A.; Lough, A.; Hopkinson, A. C.; Organ, M. G. Easily Prepared Air- and Moisture-Stable Pd-NHC (NHC = N-Heterocyclic Carbene) Complexes: A Reliable, User-Friendly, Highly Active Palladium Precatalyst for the Suzuki-Miyaura Reaction. *Chem. - A Eur. J.* **2006**, 12 (18), 4743–4748. <https://doi.org/10.1002/chem.200600251>.
  40. Someya, H.; Yorimitsu, H.; Oshima, K. Silver-Catalyzed Coupling Reactions of Alkyl Halides with Indenylolithiums. *Tetrahedron* **2010**, 66 (32), 5993–5999. <https://doi.org/10.1016/j.tet.2010.06.030>.
  41. Zhang, J.; Li, T.; Li, X.; Zhang, G.; Fang, S.; Yan, W.; Li, X.; Yang, X.; Ma, Y.; Szostak, M. An Air-Stable, Well-Defined Palladium-BIAN-NHC Chloro Dimer: A Fast-Activating, Highly Efficient Catalyst for Cross-Coupling. *Chem. Commun.* **2022**, 58 (53), 7404–7407. <https://doi.org/10.1039/d2cc02253b>.
  42. Pirkl, N.; Del Grosso, A.; Mallick, B.; Doppiu, A.; Gooßen, L. J. Dihalogen-Bridged NHC-Palladium(i) Dimers: Synthesis, Characterisation and Applications in Cross-Coupling Reactions. *Chem. Commun.* **2019**, 55 (36), 5275–5278. <https://doi.org/10.1039/c9cc02239b>.
  43. Zhou, T.; Ma, S.; Nahra, F.; Obled, A. M. C.; Poater, A.; Cavallo, L.; Cazin, C. S. J.; Nolan, S. P.; Szostak, M. [Pd(NHC)( $\mu$ -Cl)Cl]<sub>2</sub>: Versatile and Highly Reactive Complexes for Cross-Coupling Reactions That Avoid Formation of Inactive Pd(I) Off-Cycle Products. *iScience* **2020**, 23 (8), 101377. <https://doi.org/10.1016/j.isci.2020.101377>.
  44. Yang, S.; Yu, X.; Poater, A.; Cavallo, L.; Cazin, C. S. J.; Nolan, S. P.; Szostak, M. Buchwald-Hartwig Amination and C-S/S-H Metathesis of Aryl Sulfides by Selective C-S Cleavage Mediated by Air- and Moisture-Stable [Pd(NHC)( $\mu$ -Cl)Cl]<sub>2</sub>Precatalysts: Unified Mechanism for Activation of Inert C-S Bonds. *Org. Lett.* **2022**, 24 (50), 9210–9215. <https://doi.org/10.1021/acs.orglett.2c03717>.
  45. Guillet, S. G.; Voloshkin, V. A.; Saab, M.; Beliš, M.; Van Hecke, K.; Nahra, F.; Nolan, S. P. Understanding Existing and Designing Novel Synthetic Routes to Pd-PEPPSI-NHC and Pd-PEPPSI-PR3pre-Catalysts. *Chem. Commun.* **2020**, 56 (44), 5953–5956. <https://doi.org/10.1039/d0cc02262d>.
  46. Silarska, E.; Majchrzak, M.; Marciniak, B.; Trzeciak, A. M. Efficient Functionalization of Olefins by Arylsilanes Catalyzed by Palladium Anionic Complexes. *J. Mol. Catal. A Chem.* **2017**, 426, 458–464. <https://doi.org/10.1016/j.molcata.2016.07.023>.
  47. Silarska, E.; Trzeciak, A. M. Oxygen-Promoted Coupling of Arylboronic Acids with Olefins Catalyzed by [CA]<sub>2</sub>[PdX<sub>4</sub>] Complexes without a Base. *J. Mol. Catal. A Chem.* **2015**, 408, 1–11. <https://doi.org/10.1016/j.molcata.2015.07.003>.
  48. Kahan, R.; Pitak, M.; Seechurn, C. Complexes [M<sub>2</sub>X<sub>6</sub>][HPR<sub>1</sub>R<sub>2</sub>R<sub>3</sub>]<sub>2</sub>. WO2020229848A1, **2020**.
  49. Valente, C.; Çalimsiz, S.; Hoi, K. H.; Mallik, D.; Sayah, M.; Organ, M. G. The Development of Bulky Palladium NHC Complexes for the Most-Challenging Cross-Coupling Reactions. *Angew. Chemie - Int. Ed.* **2012**, 51 (14), 3314–3332. <https://doi.org/10.1002/anie.201106131>.
  50. Organ, M. G.; Çalimsiz, S.; Sayah, M.; Hoi, K. H.; Lough, A. J. Pd-PEPPSI-IPent: An Active, Sterically Demanding Cross-Coupling Catalyst and Its Application in the Synthesis of Tetra-Ortho-Substituted Biaryls. *Angew. Chemie - Int. Ed.* **2009**, 48 (13), 2383–2387. <https://doi.org/10.1002/anie.200805661>.
  51. Çalimsiz, S.; Organ, M. G. Negishi Cross-Coupling of Secondary Alkylzinc Halides with Aryl/Heteroaryl Halides Using Pd-PEPPSI-IPent. *Chem. Commun.* **2011**, 47 (18), 5181–5183. <https://doi.org/10.1039/c0cc04835f>.
  52. Atwater, B.; Chandrasoma, N.; Mitchell, D.; Rodriguez, M. J.; Organ, M. G. Pd-PEPPSI-IHeptCl: A General-Purpose, Highly Reactive Catalyst for the Selective Coupling of Secondary Alkyl Organozincs. *Chem. - A Eur. J.* **2016**, 22 (41), 14531–14534. <https://doi.org/10.1002/chem.201603603>.
  53. Atwater, B.; Chandrasoma, N.; Mitchell, D.; Rodriguez, M. J.; Pompeo, M.; Froese, R. D. J.; Organ, M. G. The Selective Cross-Coupling of Secondary Alkyl Zinc Reagents to Five-Membered-Ring Heterocycles Using Pd-PEPPSI-IHeptCl. *Angew. Chemie - Int. Ed.* **2015**, 54 (33), 9502–9506. <https://doi.org/10.1002/anie.201503941>.
  54. Pompeo, M.; Froese, R. D. J.; Hadei, N.; Organ, M. G. Pd-PEPPSI-IPentCl: A Highly Effective Catalyst for the Selective Cross-Coupling of Secondary Organozinc Reagents. *Angew. Chemie - Int. Ed.* **2012**, 51 (45), 11354–11357. <https://doi.org/10.1002/anie.201205747>.
  55. Vila, C.; Giannerini, M.; Hornillos, V.; Fañanás-Mastral, M.; Feringa, B. L. Palladium-Catalysed Direct Cross-Coupling of Secondary Alkylolithium Reagents. *Chem. Sci.* **2014**, 5 (4), 1361–1367. <https://doi.org/10.1039/c3sc53047g>.
  56. Vila, C.; Hornillos, V.; Giannerini, M.; Fañanás-Mastral, M.; Feringa, B. L. Palladium-Catalysed Direct Cross-Coupling of Organolithium Reagents with Aryl and Vinyl Triflates. *Chem. - A Eur. J.* **2014**, 20 (41), 13078–13083. <https://doi.org/10.1002/chem.201404398>.
  57. Heijnen, D.; Hornillos, V.; Corbet, B. P.; Giannerini, M.; Feringa, B. L. Palladium-Catalyzed C(Sp<sup>3</sup>)-C(Sp<sup>2</sup>) Cross-Coupling of (Trimethylsilyl)methylolithium with (Hetero)Aryl Halides. *Org. Lett.* **2015**, 17 (9), 2262–2265. <https://doi.org/10.1021/acs.orglett.5b00905>.
  58. Zhao, Q.; Meng, G.; Li, G.; Flach, C.; Mendelsohn, R.; Lalancette, R.; Szostak, R.; Szostak, M. IPr# - Highly Hindered, Broadly Applicable N-Heterocyclic Carbenes. *Chem. Sci.* **2021**, 12 (31), 10583–10589. <https://doi.org/10.1039/d1sc02619d>.
  59. Valente, C.; Baglione, S.; Candito, D.; O'Brien, C. J.; Organ, M. G. High Yielding Alkylations of Unactivated Sp<sup>3</sup> and Sp<sup>2</sup> Centres with Alkyl-9-BBN Reagents Using an NHC-Based Catalyst: Pd-PEPPSI-IPr. *Chem. Commun.* **2008**, No. 6, 735–737. <https://doi.org/10.1039/b715081d>.
  60. Peh, G. R.; Kantchev, E. A. B.; Er, J. C.; Ying, J. Y. Rational Exploration of N-Heterocyclic Carbene (NHC) Palladacycle Diversity: A Highly Active and Versatile Precatalyst for Suzuki-Miyaura Coupling Reactions of Deactivated Aryl and Alkyl Substrates. *Chem. - A Eur. J.* **2010**, 16 (13), 4010–4017. <https://doi.org/10.1002/chem.200902842>.
  61. Sun, K. X.; Zhou, J. H.; He, Q. W.; Shao, L. X.; Lu, J. M. N-Heterocyclic Carbene-Pd(II)-2-Methyl-4,5-Dihydrooxazole Complex-Catalyzed Highly Chemoselective Mono-Amination of Dichlorobenzenes. *Tetrahedron* **2020**, 76 (8), 130944. <https://doi.org/10.1016/j.tet.2020.130944>.
  62. Larock, R. C.; Yum, E. K. Synthesis of Indoles via Palladium-Catalyzed Heteroannulation of Internal Alkynes New Methodology for the Synthesis of Functionalized Indolizidine and Quinolizidine Ring Systems. *J. Am. Chem. Soc.* **1991**, 113 (9), 6689–6690.
  63. These catalysts are available at [www.sigmaaldrich.com](http://www.sigmaaldrich.com). (IPr-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub>, 939595; (SIPr-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub>, **939609**; (IPent-H)<sub>2</sub>Pd<sub>2</sub>Cl<sub>6</sub>, **939617**.

## Featured & Related Products

Description	Cat. No.
[IPr-H] <sub>2</sub> Pd <sub>2</sub> Cl <sub>6</sub>	939595
[IPent-H] <sub>2</sub> Pd <sub>2</sub> Cl <sub>6</sub>	939617
[SIPr-H] <sub>2</sub> Pd <sub>2</sub> Cl <sub>6</sub>	939609

## Related Product Categories

- Cross-coupling Catalysts
- Transition Metal Catalysis
- Buckwald Catalysts & Ligands
- NHC Ligands & Complexes
- Inorganic Catalysts
- Phosphine Ligands

## Related Technical Articles

- NHC-based Palladium Catalysts
- N-Heterocyclic Carbene (NHC) Compounds
- N-Heterocyclic Carbene (NHC) Ligands
- N-Heterocyclic Carbene-Copper Complexes
- PEPPSI™-IPent for Demanding Cross-Coupling Reactions
- Rieke® Highly Reactive Metals

### To place an order or receive technical assistance

Visit: [SigmaAldrich.com/offices](https://www.sigmaaldrich.com/offices)

For Technical Service visit: [SigmaAldrich.com.com/techservice](https://www.sigmaaldrich.com.com/techservice)

[SigmaAldrich.com](https://www.sigmaaldrich.com)

We have built a unique collection of life science brands with unrivalled experience in supporting your scientific advancements.

[Millipore®](#) [Sigma-Aldrich®](#) [Supelco®](#) [Milli-Q®](#) [SAFC®](#) [BioReliance®](#)

Merck KGaA  
Frankfurter Strasse 250  
64293 Darmstadt, Germany

