

MS5-mDLL1 Stromal Cell Line

Immortalized Cell Line

Cat. # SCC166



FOR RESEARCH USE ONLY.
NOT FOR USE IN DIAGNOSTIC PROCEDURES.
NOT FOR HUMAN OR ANIMAL CONSUMPTION.
THIS PRODUCT CONTAINS GENETICALLY MODIFIED ORGANISMS.

Pack size: $\geq 1 \times 10^6$
viable cells/vial

Store in liquid nitrogen

Data Sheet

page 1 of 3

Background:

In vivo, T-cells develop in the thymus from bone-marrow derived hematopoietic stem and progenitor cells (HSPCs). *In vitro*, various model systems have been employed to mimic thymic function supportive of T-cell development. The classic OP9-DL1 culture system utilizes a coculture of HSPC with a murine bone marrow-derived stromal cell line (OP9) that ectopically expresses the Notch ligand, Delta-like 1 (Dl1). The OP9-DL1 culture system utilizes fetal calf serum and thus contributes to variable efficiencies in T-cell differentiation and moreover suffers from low maturation rates. Similarly, three-dimensional (3D) organoid systems utilizing primary thymic tissues from murine^{1,2,3} or human⁴ sources suffer from experimental variability and difficulty in access of primary tissues.

Recently, a serum-free artificial thymic organoid (ATO) system⁵ was developed that supports highly efficient and reproducible *in vitro* differentiation and positive selection of human T-cells from multiple sources of HSPCs, including cord blood, bone marrow and peripheral blood CD34+ HSPCs. A key component of the ATO system is the MS5-mDLL1 cell line which obviates the need for primary thymic tissues. To form organoids, MS5-mDLL1 stromal cells are aggregated with HSPCs by centrifugation and cultured on a cell culture insert at the air-fluid interface. Detailed protocols for setting up the artificial thymic organoid (ATO) system are described in reference 5.

To generate the MS5-mDLL1, a murine stromal cell line (MS5)⁶ was transduced with a third-generation lentiviral vector encoding mouse DLL1 and mStrawberry.

Storage and Handling

MS5-mDLL1 cells should be stored in liquid nitrogen. The cells can be cultured for at least 10 passages after initial thawing without significantly affecting the cell marker expression and functionality.

Quality Control Testing

- Each vial contains $\geq 1 \times 10^6$ viable cells.
- Cells are negative for mycoplasma contamination.
- Cells are tested negative for infectious diseases by a Mouse Essential CLEAR panel by Charles River Animal Diagnostic Services.
- Cells are verified to be of mouse origin and negative for inter-species contamination from rat, chinese hamster, Golden Syrian hamster, human and non-human primate (NHP) as assessed by a Contamination Clear panel by Charles River Animal Diagnostic Services.

Representative Data




Figure 1. MS5-mDLL1 cells one (A) and two (B) days after thawing in a T75 flask.

References

1. Anderson G, Jenkinson EJ, Moore NC, Owen JJ (1993) MHC class II-positive epithelium and mesenchyme cells are both required for T-cell development in the thymus. *Nature* 362(6415): 70-73.
2. Plum J, De Smedt M, Defresne MP, Leclercq G, Vandekerckhove B (1994) Human CD34+ fetal liver stem cells differentiate to T cells in a mouse thymic microenvironment. *Blood* 84(5): 1587-1593.
3. Poznansky MC, et al. (2000) Efficient generation of human T cells from a tissue-engineered thymic organoid. *Nat Biotech* 18(7): 729-734.
4. Chung B, et al. (2014) Engineering the human thymic microenvironment to support thymopoiesis *in vivo*. *Stem Cells* 32(9): 2386-2396.
5. Seet CS, et al. (2017) Generation of mature T cells from human hematopoietic stem and progenitor cells in artificial thymic organoids. *Nat Methods* 14(5): 521-530.
6. Itoh K, et al. (1989) Reproducible establishment of hematopoietic supportive stromal cell lines from murine bone marrow. *Exp Hematol* 17(2): 145-153.

Please visit www.millipore.com for additional product information and references.

Submit your published journal article, and earn credit toward future purchases. Visit www.millipore.com/publicationrewards to learn more!

Protocols

Thawing Cells

1. Do not thaw the cells until the recommended medium is on hand. Cells can grow on normal tissue cultureware surfaces without any additional coating.

Cells are thawed and expanded in High Glucose DMEM (Sigma Cat. No. D6429), 10% FBS (Cat. No. ES-009-B), and 1X Penicillin-Streptomycin Solution (Cat. No. TMS-AB2-C) (optional).

2. Remove the vial of frozen MS5-mDLL1 cells from liquid nitrogen and incubate in a 37°C water bath. Closely monitor until the cells are completely thawed. Maximum cell viability is dependent on the rapid and complete thawing of frozen cells.

IMPORTANT: Do not vortex the cells.

3. As soon as the cells are completely thawed, disinfect the outside of the vial with 70% ethanol. Proceed immediately to the next step.

4. In a laminar flow hood, use a 1 or 2 mL pipette to transfer the cells to a sterile 15 mL conical tube. Be careful not to introduce any bubbles during the transfer process.

5. Using a 10 mL pipette, slowly add dropwise 9 mL of MS5-DLL1 Expansion Medium (Step 1 above) to the 15 mL conical tube.

IMPORTANT: Do not add the entire volume of media all at once to the cells. This may result in decreased cell viability due to osmotic shock.

6. Gently mix the cell suspension by slowly pipetting up and down twice. Be careful not to introduce any bubbles.

IMPORTANT: Do not vortex the cells.

7. Centrifuge the tube at 300 x g for 2-3 minutes to pellet the cells.

8. Decant as much of the supernatant as possible. Steps 5-8 are necessary to remove residual cryopreservative (DMSO).

9. Resuspend the cells in 45 – 50 mL of MS5-DLL1 Expansion Medium.

10. Transfer the cell mixture to a **T175** tissue culture flask.

Note: *MS5-mDLL1 cells proliferate rapidly.*

11. Incubate the cells at 37°C in a humidified incubator with 5% CO₂.

Subculturing Cells

1. Carefully remove the medium from the T175 tissue culture flask containing the confluent layer of MS5-mDLL1 cells.

2. Rinse the T175 flask twice with 15-20 mL 1X PBS. Aspirate after each rinse.

3. Apply 10-15 mL of Accutase or trypsin-EDTA solution and incubate in a 37°C incubator for 3-5 minutes.

4. Inspect the flask and ensure the complete detachment of cells by gently tapping the side of the flask with the palm of your hand.

5. Add 12 mL of MS5-DLL1 Expansion Medium to the plate.

6. Gently rotate the flask to mix the cell suspension. Transfer the dissociated cells to a 50 mL conical tube.

7. Centrifuge the tube at 300 x g for 3-5 minutes to pellet the cells.

8. Discard the supernatant, then loosen the cell pellet by tapping the tip of the tube with a finger.

9. Apply 2-5 mL of MS5-DLL1 Expansion Medium to the conical tube and resuspend the cells thoroughly.

IMPORTANT: Do not vortex the cells.

10. Count the number of cells using a hemocytometer.

11. Plate the cells to the desired density (typical split ratio is 1:6- 1:9). Cells proliferate extremely rapidly.

Cryopreservation of Cells

MS5-mDLL1 mouse bone marrow stromal cell line transduced with human DLL1 may be frozen in the expansion medium plus 10% DMSO using a Nalgene slow freeze Mr. Frosty container.

█ antibodies █ Multiplex products █ biotools █ cell culture █ enzymes █ kits █ proteins/peptides █ siRNA/cDNA products

Please visit www.millipore.com for additional product information, test data and references

EMD Millipore Corporation, 28820 Single Oak Drive, Temecula, CA 92590, USA 1-800-437-7500

Technical Support: T: 1-800-MILLIPORE (1-800-645-5476) • F: 1-800-437-7502

FOR RESEARCH USE ONLY. Not for use in diagnostic procedures. Not for human or animal consumption. Purchase of this Product does not include any right to resell or transfer, either as a stand-alone product or as a component of another product. Any use of this Product for purposes other than research is strictly prohibited.

EMD Millipore®, the M mark, Upstate®, Chemicon®, Linco® and all other registered trademarks, unless specifically identified above in the text as belonging to a third party, are owned by Merck KGaA, Darmstadt, Germany. Copyright ©2008-2018 Merck KGaA, Darmstadt, Germany. All rights reserved.



We Buy 100% Certified
Renewable Energy

ACADEMIC USE AGREEMENT
(subject to local law)

THIS PRODUCT MAY ONLY BE USED BY INDIVIDUALS EMPLOYED BY AN ACADEMIC INSTITUTION AND IS INTENDED SOLELY TO BE USED FOR ACADEMIC RESEARCH, WHICH IS FURTHER DEFINED BELOW. BY OPENING THIS PRODUCT, YOU ("PURCHASER") HEREBY REPRESENT THAT YOU HAVE THE RIGHT AND AUTHORITY TO LEGALLY BIND YOURSELF AND/OR YOUR EMPLOYER INSTITUTION, AS APPLICABLE, AND CONSENT TO BE LEGALLY BOUND BY THE TERMS OF THIS ACADEMIC USE AGREEMENT. IF YOU DO NOT AGREE TO COMPLY WITH THESE TERMS, YOU MAY NOT OPEN OR USE THE PRODUCT AND YOU MUST CALL MILLIPORESIGMA ("SELLER") CUSTOMER SERVICE (1-800-645-5476) TO ARRANGE TO RETURN THE PRODUCT FOR A REFUND.

"Product" means MS5-mDLL1 Stromal Cell Line (SCC166)

"Academic Research" means any internal *in vitro* research use by individuals employed by an academic institution. Academic Research specifically excludes the following uses of whatever kind or nature:

- Re-engineering or copying the Product
- Making derivatives, modifications, or functional equivalents of the Product
- Obtaining patents or other intellectual property rights claiming use of the Product
- Using the Product in the development, testing, or manufacture of a Commercial Product
- Using the Product as a component of a Commercial Product
- Reselling or licensing the Product
- Using the Product in clinical or therapeutic applications including producing materials for clinical trials
- Administering the Product to humans
- Using the Product in collaboration with a commercial or non-academic entity

"Commercial Product" means any product intended for: (i) current or future sale; (ii) use in a fee-for-service; or (iii) any diagnostic, clinical, or therapeutic use.

Access to the Product is limited solely to those officers, employees, and students of PURCHASER's academic institution who need access to the Product to perform Academic Research. PURCHASER shall comply with all applicable laws in its use and handling of the Product and shall keep it under reasonably safe and secure conditions to prevent unauthorized use or access.

These use restrictions will remain in effect for as long as PURCHASER possesses the Product.

COMMERCIAL OR NON-ACADEMIC ENTITIES INTERESTED IN PURCHASING OR USING THE PRODUCT MUST CONTACT licensing@emdmillipore.com AND AGREE TO SEPARATE TERMS OF USE PRIOR TO USE OR PURCHASE.

GMO

This product contains genetically modified organisms.
Este producto contiene organismos genéticamente modificados.
Questo prodotto contiene degli organismi geneticamente modificati.
Dieses Produkt enthält genetisch modifizierte Organismen.
Ce produit contient des organismes génétiquement modifiés.
Dit product bevat genetisch gewijzigde organismen.
Tämä tuote sisältää geneettisesti muutettuja organismeja.
Denna produkt innehåller genetiskt ändrade organismer.

█ antibodies █ Multiplex products █ biotools █ cell culture █ enzymes █ kits █ proteins/peptides █ siRNA/cDNA products

Please visit www.millipore.com for additional product information, test data and references

EMD Millipore Corporation, 28820 Single Oak Drive, Temecula, CA 92590, USA 1-800-437-7500

Technical Support: T: 1-800-MILLIPORE (1-800-645-5476) • F: 1-800-437-7502

FOR RESEARCH USE ONLY. Not for use in diagnostic procedures. Not for human or animal consumption. Purchase of this Product does not include any right to resell or transfer, either as a stand-alone product or as a component of another product. Any use of this Product for purposes other than research is strictly prohibited.

EMD Millipore®, the M mark, Upstate®, Chemicon®, Linco® and all other registered trademarks, unless specifically identified above in the text as belonging to a third party, are owned by Merck KGaA, Darmstadt, Germany. Copyright ©2008-2018 Merck KGaA, Darmstadt, Germany. All rights reserved.



We Buy 100% Certified
Renewable Energy