

59808 Atto 740 NHS ester

Application


Atto 740 belongs to a new generation of fluorescent labels for the near infrared spectral region. The dye is designed for application in the area of life science, e.g. labeling of DNA, RNA or proteins. Characteristic features of the dye are strong absorption and good fluorescence as well as excellent thermal and photo-stability.

Atto 740 is a cationic dye. After coupling to a substrate the dye carries a net electrical charge of +1. Atto 740 is a pH sensitive product. While practically stable up to pH 7.4 (PBS-buffer), it slowly degrades at higher pH. If exposed to higher pH for coupling purposes, we recommend reducing the pH immediately after completion of the reaction.

Product Description

MW	665 g/mol
λ_{abs}	743 nm
ϵ_{max}	$1.2 \times 10^5 \text{ M}^{-1} \text{ cm}^{-1}$
λ_{fl}	763 nm
η_{fl}	10 %
τ_{fl}	0.6 ns
CF ₂₆₀	0.07
CF ₂₈₀	0.07

Optical data of the carboxy derivative (in aqueous solution)



Directions for labeling of proteins with Atto 740 NHS ester

Dissolve protein in of bicarbonate buffer (0.1 M, preferably of pH 8.3) at 2 mg/ml. Concentrations below 2 mg protein per ml will decrease labeling efficiency. Protein or peptide solutions must be free of any amine-containing substances such as Tris, glycine or ammonium salts. Antibodies in solutions of Tris buffers may be dialyzed against 10-20 mM PBS. The desired pH can be obtained by adding 0.1 ml of 1 M sodium bicarbonate buffer, pH 8.3, for each ml of dialyzed antibody solution.

Dissolve Atto 740 NHS ester in amine-free, dry DMF or DMSO at 2 mg/ml (e.g. 1 mg Atto 740 NHS in 500 μl). This solution should always be prepared immediately before conjugation.

As number and position of amine groups vary between different proteins, the optimum of dye/protein ratio also varies. Wherefore we recommend, to try out different ratios when labeling a certain protein for the first time.



In general a molar ratio of 1-2 may be suitable for smaller proteins, much higher ratios for large proteins. To obtain a ratio in this range, add a twofold molar excess of reactive dye to the protein solution. For an antibody, a ratio of 8 or higher may be a good starting point. For such ratio, add 50 µl of dye solution to 1 ml protein solution. Variations due to different reactivities of both the protein (number, position and reactivity of amine groups) may occur. This may necessitate individual optimization of the dye to protein ratio used in the reaction in order to optimize the results.

Incubate the reaction at room temperature for 30 to 60 min under constant or repeated stirring.

Separation of labeled proteins

The labeled protein can be separated from unreacted dye by gel permeation chromatography, e.g. using a Sephadex™ G-25, G50 or Bio-Gel™ P-10 column. We recommend to use Sephadex™ G-25. The column should have a diameter of at least 1 cm and a length of 12 cm. It can be equilibrated with phosphate buffer of pH 7.2 (22 mM) or another buffer of choice. The same buffer can be used for elution. Usually, the first fluorescent band is the labelled protein, while free dye will elute in a second fluorescent band.

In case you have to work with diluted samples, you may purify the conjugate by extensive dialysis. But this is less efficient and not as fast as purification by gel filtration.

Storage of conjugates

In general, conjugates should be stored under the same conditions used for the unlabeled protein. For storage in solution at 4°C, 2 mM sodium azide can be added as a preservative. Typically, protein conjugates will be stable for several months. For long-term storage, aliquots may be frozen at -20°C to avoid repeated freezing and thawing. Protect from light. If your protein tends to instability please use one of our BioStab solutions specially designed for stabilization of proteins.

After long-term storage of conjugate solutions we recommend to centrifuge in a micro-centrifuge before use. This will remove any aggregates which might have formed.

Directions for oligo-nucleotides with Atto 740 NHS ester

Prepare a solution of 0.1 mM solution (e.g. 5 nmol in 50 µl) of amino-modified oligo-nucleotide in carbonate buffer (0.2 M, pH 8-9). Prepare a solution of 5 mg/ml activated label in anhydrous DMF. Add ~ 50 µl of oligo-nucleotide solution to 30 µl of label solution. Incubate the reaction at room temperature for 2 hours under shaking. If longer reaction times are required, the pH value should be diminished to pH 7-7.5.

Separation of labeled oligo-nucleotides

Conjugated oligo-nucleotides can be separated from free dye using gel filtration or reversed phase HPLC.

Sephadex is a registered trademark of GE Healthcare

Storage: Store at -20°C and protected from light.

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

The vibrant M, Supelco, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. Detailed information on trademarks is available via publicly accessible resources.
© 2018 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the US and Canada.

