

Technical Bulletin

Steroids Standard Mixture

Catalogue number SMB00968

Product Description

Steroid hormones are essential in numerous physiological processes including metabolism, homeostasis, development, growth, reproduction, and energy metabolism. They also have profound roles in molecular and cellular mechanisms.

Thus, an accurate quantitative or qualitative analysis is required for research purposes. This can be achieved by using stable labeled steroid standards.

A method of choice for target or untargeted metabolomics analysis of different classes of steroid's is Liquid Chromatography – Mass Spectrometry (LC-MS). For targeted or untargeted metabolomics overcoming ion suppression, ion variability and drifting signal are challenging aspects of analysis. Use of a deuterated steroids mixture could overcome the challenges mentioned above.

The Steroids Standard Mixture is dissolved in methanol. This mixture is formulated for use as a concentration standard for high-throughput LC/MS metabolomic analysis.

- All deuterated steroids are dissolved in methanol and ready to use. Total vial volume is 1 mL.
- This mixture can be used for targeted and untargeted approaches.
- Ideal for LC-MS studies as internal standards.

Components of the deuterated Steroids Standard Mixture

* See batch-specific CoA for measured concentrations

Components
Aldosterone-d7
11-Deoxycortisol-d5
Androstenedione-c3
Pregnenolone-c2d2
Corticosterone-d4
Allopregnanolone-d5
Etiocolanolone-d5
Hydrocortisone sulfate-d4
Cortisone-d8
Testosteron-d3
Cortisol-d4
5α-Dihydrotestosteron-d3
Progesterone-d9
17-OH Progesterone-d8
Dehydroepiandrosterone-d6 sulfate sodium salt

Precautions and Disclaimer

This product is for R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

Store the sealed vials at -20 °C.

Preparation Instructions

The Steroids Standard Mixture is a ready-to-use solution.

Procedure

1. To 50 μ L plasma Samples add 6.25 μ L of deuterated Steroid Standard Mixture.
2. Add 10.0 μ L of antioxidant (butylated hydroxytoluene and ethylenediaminetetraacetic acid (EDTA)) and 10 μ L of 1000 nM 1-cyclohexyluriedo-3-dodecanoic acid and 1-phenyl-3-hexadecanoic acid urea.
3. Dilute to final volume of 250 μ L with 1:1 acetonitrile: methanol.
4. Vortex and incubate at 20 °C for 30 minutes to precipitate protein, and centrifugate at 15,000 rcf for 5 minutes.
5. Pass through a 0.2 μ m PVDF filter plate and store at -20 °C until analysis.

Reagents required and not provided

Butylated Hydroxytoluene – B1378; EDTA – EDS; 1-cyclohexyluriedo-3-dodecanoic acid (CUDA) and 1-Phenyl 3-Hexadecanoic Acid Urea (PHAU); acetonitrile (ACN) - 34851; Methanol (CH₃OH – 34860.

References

Barupal, D.K, *et al.*, "A Comprehensive Plasma Metabolomics Dataset for a Cohort of Mouse Knockouts within the International Mouse Phenotyping Consortium." *Metabolites*, **9**, 101 (2019).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Terms and Conditions of Sale

Warranty, use restrictions, and other conditions of sale may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck operates
as MilliporeSigma in the U.S. and Canada.

Merck and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2022 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

smb00968pis Rev 03/23

