

## Product Information

## $\beta$ -Glucuronidase from *Helix pomatia*

Type H-1, partially purified powder,  $\geq 300,000$  units/g solid

**G0751**

### Product Description


CAS Registry Number: 9001-45-0

Enzyme Commission (EC) Number: 3.2.1.31

Synonyms:  $\beta$ -D-Glucuronide glucuronosohydrolase

Glucuronidation, or conjugation with glucuronic acid, by the human UDP-glucuronosyltransferase (UGT) family of enzymes plays an important role in the metabolic fate of many drugs and other xenobiotics. This biosynthetic reaction also has a role in the conjugation and excretion of endogenous substrates, such as steroids, bilirubin, and bile acids.<sup>1</sup> UGT activity results in the conjugation of glucuronic acid to substrates that contain sulfhydryl, hydroxyl, aromatic amino, or carboxylic acid moieties. The resulting glucuronides are more polar (water-soluble) than the parent organic substrate and are generally excreted through the kidney.

$\beta$ -glucuronidase catalyzes the general reaction:



$\beta$ -Glucuronidase Type H-1 has been used for the enzymatic hydrolysis of these metabolites from urine,<sup>2-4</sup> plasma,<sup>5</sup> serum,<sup>6,7</sup> and bile<sup>8</sup> prior to analysis by various means. The enzyme has also been used for the digestion of fungal mycelium.<sup>9</sup> Typically, between 1-20 units of glucuronidase are used per  $\mu\text{L}$  of plasma, urine, or bile for the enzymatic hydrolysis of glucuronides present in these samples.<sup>2-8</sup> The exact amount needed will depend on the specific conditions used and must be determined empirically.

$\beta$ -Glucuronidase Type H-1 from *Helix pomatia* is a partially purified, essentially salt-free powder of enzymes derived from the Roman snail. Many  $\beta$ -glucuronidases derived from mollusks also contain sulfatase activity. For this reason, sulfatase activity is also determined.

Several theses<sup>10</sup> and dissertations<sup>11-18</sup> have cited use of product G0751 in their protocols.

### Optimal pH

- Glucuronidase activity: 4.5 to 5.0
- Sulfatase activity:  $\sim 6.2$

### Inhibitors

- D-glucuronic acid (Cat. No. G5269)
- D-galacturonic acid (Cat. No. 48280)
- D-glucaro-1,4-lactone

### Substrates

- 5-Bromo-6-chloro-3-indolyl  $\beta$ -D-glucuronide (Cat. No. B4532)
- 6-Bromo-2-naphthyl  $\beta$ -D-glucuronide (Cat. No. B7877)
- 5-Bromo-4-chloro-3-indolyl  $\beta$ -D-glucuronide sodium salt tablet (Cat. No. B8174)
- 8-Hydroxyquinoline glucuronide sodium salt (Cat. No. 38153)
- 4-Methylumbelliferyl  $\beta$ -D-glucuronide (Cat. No. M9130)
- 4-Nitrophenyl  $\beta$ -D-glucuronide (Cat. Nos. N1627, 73677)

### Glucuronidase Activity

$\geq 300,000$  units per gram solid

Unit Definition: One Sigma or modified Fishman unit will liberate 1.0  $\mu\text{g}$  of phenolphthalein from phenolphthalein glucuronide per hour at 37 °C at pH 5.0 (30-minute assay).

### Sulfatase Activity

$\geq 10,000$  units per gram solid

Unit Definition: One unit of sulfatase will hydrolyze 1.0  $\mu\text{mole}$  of *p*-nitrocatechol sulfate per hour at pH 5.0 at 37 °C.

**MERCK**

## Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

## Storage/Stability

Store the product at -20 °C. When stored at -20 °C, the enzyme retains activity for at least 3 years.

## References

1. Tephly, T.R. *et al.*, *Adv. Pharmacol.*, **42**, 343-346 (1998).
2. Vogt, B. *et al.*, *Nephrol. Dial. Transplant.*, **17(5)**, 753-758 (2002).
3. LeBeau, M.A. *et al.*, *J. Forensic Sci.*, **45(5)**, 1133-1141 (2000).
4. Santos, A. *et al.*, *Clin. Cancer Res.*, **6(5)**, 2012-2020 (2000).
5. Sugimoto, T. *et al.*, *Xenobiotica*, **29(8)**, 803-814 (1999).
6. Dalu, A. *et al.*, *Prostate*, **37(1)**, 36-43 (1998).
7. Lamartiniere, C.A. *et al.*, *Tox. Sci.*, **65(2)**, 228-238 (2002).
8. Pozzi, E.J.S. *et al.*, *J. Pharmacol. Exp. Ther.*, **306(1)**, 279-286 (2003).
9. Thevissen, K. *et al.*, *J. Biol. Chem.*, **272(51)**, 32176-32181 (1997).
10. Zamzow, Emily, "Measuring Urinary Hydroxylated PAH Metabolites as Biomarkers of Exposure to Diesel Exhaust in Underground Miners". University of Washington, M.S. thesis, p. A.16 (2017).
11. Christoffel, Julie, "Endocrine potential of Genistein, Resveratrol and 8-Prenylnaringenin in gonadal and thyroid axes and related organs". Christian-Albrechts-Universität zu Kiel, Ph.D. dissertation, p. 40 (2004).
12. Larkin, Theresa Ann, "Soy Isoflavone bioavailability - effects of probiotic and prebiotic consumption and oil supplementation". University of Wollongong, Ph.D. dissertation, p. 49 (2005).
13. Aboukhaddour, Reem, "*Pyrenophora tritici-repentis*: Genome Structural Organization and Genetic Diversity". University of Manitoba, Ph.D. dissertation, p. 29 (2008).

14. Graham, Lisa Anne, "Environmental Estrogens: Assessing Human Gestational Exposure and Interactions with the Estrogen Receptor". University of Canterbury, Ph.D. dissertation, p. 147 (2012).
15. Song, Brian Jay, "The chemistry of flavonoids in model beverages and human milk". Purdue University, Ph.D. dissertation, pp. 66, 87 (2013).
16. Yoo, Hong Sik, "Effects of inter-strain differences in the metabolism of trichloroethylene on liver and kidney toxicity". University of North Carolina at Chapel Hill, Ph.D. dissertation, pp. 29, 60 (2014).
17. Luo, Yu-Syuan, "Inter-individual variability in metabolism and toxicity of trichloroethylene and tetrachloroethylene". Texas A&M University, Ph.D. dissertation, p. 130 (2018).
18. Soundararajan, Avinash, "Novel clinico-molecular insights into the role of bisphenol-A (BPA) in the etiology of diabetes". Deakin University, Ph.D. dissertation, p. 43 (2018).

## Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

## Technical Assistance

Visit the tech service page at [SigmaAldrich.com/techservice](https://www.sigmaaldrich.com/techservice).

## Standard Warranty

The applicable warranty for the products listed in this publication may be found at [SigmaAldrich.com/terms](https://www.sigmaaldrich.com/terms).

## Contact Information

For the location of the office nearest you, go to [SigmaAldrich.com/offices](https://www.sigmaaldrich.com/offices).

The life science business of Merck operates as MilliporeSigma in the U.S. and Canada.

Merck and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2022 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.  
G0751dat Rev 04/22 AC,GCY,MAM

