3050 Spruce Street, St. Louis, MO 63103 USA Tel: (800) 521-8956 (314) 771-5765 Fax: (800) 325-5052 (314) 771-5757 email: techservice@sial.com sigma-aldrich.com

# **Product Information**

### **Anti-Oxytocin Receptor**

produced in rabbit, affinity isolated antibody

Catalog Number O4389

#### **Product Description**

Anti-Oxytocin Receptor is produced in rabbit using as immunogen a synthetic peptide conjugated to KLH. The peptide corresponds to the N-terminal extracellular domain of human oxytocin receptor. The antibody is affinity-purified using the immunizing peptide immobilized on agarose.

Anti-Oxytocin Receptor specifically recognizes human oxytocin receptor by immunohistochemistry with formalin-fixed, paraffin-embedded tissues. Not tested for other uses. The immunizing peptide has 95% homology with the rat and mouse gene. Other species reactivity has not been confirmed.

Oxytocin Receptor has been reported in breast, central nervous system, heart, ovary, placenta, prostate, salivary gland, testis, uterus, and vessel. ESTs (Expressed Sequence Tags) have been isolated from connective tissue, embryo, heart, lung, and melanocyte/uterus/fetal heart, skin, and uterus libraries.

#### Reagent

Supplied as a solution of 1 mg/ml in phosphate buffered saline, pH 7.7, containing 0.1% sodium azide as a preservative.

#### **Precautions and Disclaimer**

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

## Storage/Stability

For continuous use, store at 2-8 °C for up to one month. For extended storage, freeze in working aliquots. Repeated freezing and thawing, or storage in "frost-free" freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilution samples should be discarded if not used within 12 hours.

#### **Product Profile**

 $\frac{Immunohistochemistry}{concentration of 9 \ \mu g/ml} \ is \ determined \ using \ human prostate, \ glandular \ epithelium.$ 

**Note:** In order to obtain the best results and assay sensitivity in different techniques and preparations, we recommend determining optimal working dilutions by titration test.

This product manufactured by MBL International.

CS,KAA,PHC 11/09-1