

67455 Atto 514 NHS ester


Application

Atto 514 is a new hydrophilic fluorescent label with excellent water solubility. The dye exhibits strong absorption, high fluorescence quantum yield and exceptional thermal and photo-stability. Thus Atto 514 is highly suitable for single-molecule detection applications and high-resolution microscopy such as PALM, dSTORM, STED etc. Additionally the dye highly qualifies to be applied in flow cytometry (FACS), fluorescence in-situ hybridization (FISH) and many more. The fluorescence is excited most efficiently in the range 510 - 535 nm. A suitable source of excitation is the 514 nm line of the Argon-Ion laser.

Product Description

MW	1111 g/mol
λ_{abs}	511 nm
ϵ_{max}	$1.15 \times 10^5 \text{ M}^{-1} \text{ cm}^{-1}$
λ_{fl}	532 nm
η_{fl}	85 %
τ_{fl}	3.9 ns
CF ₂₆₀	0.21
CF ₂₈₀	0.07

Optical data of the carboxy derivative (in aqueous solution)



Directions for labeling of proteins with Atto 514 NHS ester

Dissolve protein in of bicarbonate buffer (0.1 M, preferably of pH 8.3) at 2 mg/ml. Concentrations below 2 mg protein per ml will decrease labeling efficiency. Protein or peptide solutions must be free of any amine-containing substances such as Tris, glycine or ammonium salts. Antibodies in solutions of Tris buffers may be dialyzed against 10-20 mM PBS. The desired pH can be obtained by adding 0.1 ml of 1 M sodium bicarbonate buffer, pH 7.5, for each ml of dialyzed antibody solution.

Dissolve Atto 514 NHS ester in amine-free, dry DMF or DMSO at 2 mg/ml (e.g. 1 mg Atto 514 NHS in 500 µl). This solution should always be prepared immediately before conjugation.

As number and position of amine groups vary between different proteins, the optimum of dye/protein ratio also varies. Wherefore we recommend to try out different ratios when labeling a certain protein for the first time.

In general a ratio of 1-2 may be suitable. To obtain a ratio in this range, add a twofold molar excess of reactive dye to the protein solution. For an antibody, add 10 µl of dye solution to 1 ml protein solution. Incubate the reaction at room temperature for 30 to 60 min under constant or repeated stirring.

Separation of labeled proteins

The labeled protein can be separated from unreacted dye by gel permeation chromatography, e.g. using a Sephadex™ G-25, G50 or Bio-Gel™ P-10 column. We recommend to use Sephadex™ G-25. The column should have a diameter of at least 1 cm and a length of 12 cm. It can be equilibrated with phosphate buffer of pH 7.2 (22 mM) or another buffer of choice. The same buffer can be used for elution. Usually, the first fluorescent band is the labelled protein, while free dye will elute in a second fluorescent band.

In case you have to work with diluted samples, you may purify the conjugate by extensive dialysis. But this is less efficient and not as fast as purification by gel filtration.

Storage of conjugates

In general, conjugates should be stored under the same conditions used for the unlabeled protein. For storage in solution at 4°C, 2 mM sodium azide can be added as a preservative. Typically, protein conjugates will be stable for several months. For long-term storage, aliquots may be frozen at -20°C to avoid repeated freezing and thawing. Protect from light. If your protein tends to instability please use one of our BioStab solutions specially designed for stabilization of proteins.

After long-term storage of conjugate solutions we recommend to centrifuge in a micro-centrifuge before use. This will remove any aggregates which might have formed.

Directions for oligo-nucleotides with Atto 514 NHS ester

Prepare a solution of 0.1 mM solution (e.g. 5 nmol in 50 µl) of amino-modified oligo-nucleotide in carbonate buffer (0.2 M, pH 8-9). Prepare a solution of 5 mg/ml activated label in anhydrous DMF. Add ~50 µl of oligo-nucleotide solution to 30 µl of label solution. Incubate the reaction at room temperature for 2 hours under shaking. If longer reaction times are required, the pH value should be diminished to pH 7-7.5.

Separation of labeled oligo-nucleotides

Conjugated oligo-nucleotides can be separated from free dye using gel filtration or reversed phase HPLC.

Storage: protected from light at -20°C

Sephadex is a registered trademark of GE Healthcare

Precautions and Disclaimer:

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

The vibrant M and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates.

Detailed information on trademarks is available via publicly accessible resources.

© 2018 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the US and Canada.

